

The University of Arkansas for Medical Sciences
Trauma Clinical Practice Management Guideline

SUBJECT: Solid Organ Injury Management Guideline (Spleen, Liver, Kidney)

AUTHORS: Jordan Greer, MD, PharmD; Kyle J. Kalkwarf, MD

APPROVED: 12//2018

UPDATED: 11/26/2025

EFFECTIVE: 11/26/2025

PURPOSE: To determine when non-operative management of liver, kidney, and spleen injuries is indicated.

DEFINITIONS:

SPLEEN INJURIES:

- Grade I Subcapsular hematoma, <10% surface area
 Parenchymal laceration < 1 cm in depth
 Capsular tear
- Grade II Subcapsular hematoma, 10-50% surface area
 Intraparenchymal hematoma <5 cm in diameter
 Parenchymal laceration 1-3 cm in depth
- Grade III Subcapsular hematoma, >50% surface area or expanding
 Ruptured subcapsular or intraparenchymal hematoma \geq 5 cm
 Parenchymal laceration >3 cm in depth
- Grade IV Any injury in the presence of a splenic vascular injury or active bleeding confined within
 the splenic capsule.
 Parenchymal laceration involving segmental or hilar vessels producing >25%
 devascularization
- Grade V Any injury in the presence of a splenic vascular injury with active bleeding extends
 beyond the spleen into the peritoneum.
 Shattered spleen

LIVER INJURIES:

- Grade I Subcapsular hematoma <10% surface area
 Parenchymal laceration <1 cm in depth
- Grade II Subcapsular hematoma 10-50% surface area
 Intraparenchymal hematoma <10 cm in diameter
 Parenchymal laceration 1-3 cm in depth and \leq 10 cm in length
- Grade III Subcapsular hematoma >50% surface area or expanding
 Ruptured subcapsular or parenchymal hematoma
 Intraparenchymal hematoma >10 cm
 Parenchymal laceration >3 cm in depth
 Any injury in the presence of a liver vascular injury or active bleeding contained within
 the liver parenchyma
- Grade IV Parenchymal disruption involving 25-75% of a hepatic lobe
 Active bleeding extending beyond the liver parenchyma into the peritoneum

These guidelines were prepared by the UAMS Trauma/EGS Service. They are intended to serve only as a guideline based on current review of the medical literature and practice. They are neither policies nor protocols. Their use is at the discretion of the managing physician.

The University of Arkansas for Medical Sciences
Trauma Clinical Practice Management Guideline

Grade V Parenchymal disruption involving >75% of the hepatic lobe
Juxtahepatic venous injuries to include retrohepatic vena cava and/or central major hepatic veins

RENAL INJURIES:

Grade I Subcapsular hematoma and/or parenchymal contusion without laceration

Grade II Perirenal hematoma confined to Gerota's fascia

Parenchymal laceration ≤ 1 cm in depth without urinary extravasation

Grade III Parenchymal laceration >1 cm in depth without collecting system rupture or urinary extravasation

Any injury in the presence of a kidney vascular injury or active bleeding contained within Gerota's fascia

Grade IV Parenchymal laceration extending into the urinary collecting system with urinary extravasation

Renal pelvis laceration and/or complete ureteropelvic disruption

Segmental renal vein or artery injury

Active bleeding beyond Gerota's fascia into the retroperitoneum or peritoneum

Segmental or complete kidney infarction(s) due to vessel thrombosis without active bleeding

Grade V Main renal artery or vein laceration or avulsion of hilum

Devascularized kidney with active bleeding

Shattered kidney with loss of identifiable parenchymal renal anatomy

CRITERIA:

1. Splenic Injuries

A. Non-operative management of splenic injuries can be considered when all of the following conditions have been met:

1. Diagnosis of injury on CT scan/FAST
2. Hemodynamic stability
3. Grade 1-4 injury
4. No other major intra-abdominal injury
5. No other major sources of blood loss
6. Available for monitoring except for short operative procedures
7. No other pre-morbid illnesses that suggest the patient could not tolerate blood loss (e.g., severe ischemic heart disease)
8. Willingness to receive a blood transfusion
9. Consider intervention for those with moderate-to-severe TBI to avoid hypotension

B. All patients with Grade 4-5 injuries should be evaluated for an urgent or emergent angiogram with embolization within 1 hour of presentation.

C. Admit all Grade 4-5 injuries to the SICU for close hemodynamic monitoring.

D. Consider repeating the CTA in 48-72 hours after injury to evaluate for a

This guideline was prepared by the UAMS ACS Division based on a recent literature review. It is intended to support clinical decision-making and should be used at the discretion of the trauma team, as it is not a fixed policy or protocol.

The University of Arkansas for Medical Sciences
Trauma Clinical Practice Management Guideline

pseudoaneurysm.

- E. UAMS Trauma Registry data indicate patients >60 are at high risk for failed non-operative management.
- F. **Immunizations:** Please refer to the “Post-Splenectomy Vaccines” Guideline for proper management.
- G. **Thromboprophylaxis:** Please refer to the “Venous Thromboembolism Prophylaxis (DVT/PE)” Guideline for management.

2. Liver Injuries

- A. Non-operative management of liver injuries can be considered when all of the following conditions have been met:
 - 1. Liver injury diagnosed on CT scan with normalizing vital signs, Grade I to IV
 - 2. Injury not into the hilum
 - 3. A rim of blood around the liver
- B. Admit all Grade 4 or higher liver lacerations to the ICU
 - 1. Monitor hourly vital signs
 - 2. Bedrest
 - 3. NPO
 - 4. Serial Hgb/Hct q 6 hours until stable
- C. Bile Leaks
 - 1. If <300ml/day at time of diagnosis, then observe or manage with percutaneous drainage alone for up to 14 days.
 - 2. If >300ml/day at the time of diagnosis, proceed with ERCP and sphincterotomy, or if the above treatment fails to resolve the leak within 14 days.
- D. Consider repeating the CTA in 48-72 hours from the injury to evaluate for a pseudoaneurysm.
- E. **Thromboprophylaxis:** Please refer to the “Venous Thromboembolism Prophylaxis (DVT/PE)” Guideline for management.

3. Renal Injuries

- A. Non-operative management of kidney injuries can be considered when all the following conditions have been met:
 - 1. No other indication for exploration
 - 2. Any grade in a hemodynamically stable patient
- B. Consider admitting all grade 4 or higher kidney injuries to the ICU

REFERENCES:

1. Stassen NA, Bhullar I, Cheng JD, Crandall ML, Friese RS, Guillamondegui OD, Jawa RS, Maung AA, Rohs Jr TJ, Sangosanya A, Schuster KM. Selective nonoperative management of blunt splenic injury: an Eastern Association for the Surgery of Trauma practice management guideline. *Journal of Trauma and Acute Care Surgery*. 2012 Nov 1;73(5):S294-300.
2. Kozar RA, Crandall M, Shanmuganathan K, Zarzaur BL, Coburn M, Cribari C, Kaups K, Schuster K, Tominaga GT; AAST Patient Assessment Committee. Organ injury scaling 2018 update: Spleen, liver, and kidney. *J Trauma Acute Care Surg*. 2018 Dec;85(6):1119-1122.

This guideline was prepared by the UAMS ACS Division based on a recent literature review. It is intended to support clinical decision-making and should be used at the discretion of the trauma team, as it is not a fixed policy or protocol.

The University of Arkansas for Medical Sciences
Trauma Clinical Practice Management Guideline

3. Crichton JCI, Naidoo K, Yet B, Brundage SI, Perkins Z. The role of splenic angioembolization as an adjunct to nonoperative management of blunt splenic injuries: A systematic review and meta-analysis. *J Trauma Acute Care Surg.* 2017 Nov;83(5):934-943.
4. Chehab M, Afaneh A, Bible L, Castanon L, Hanna K, Ditillo M, Khurram M, Asmar S, Joseph B. Angioembolization in intra-abdominal solid organ injury: Does delay in angioembolization affect outcomes? *J Trauma Acute Care Surg.* 2020 Oct;89(4):723-729.
5. Scarborough JE, Ingraham AM, Liepert AE, Jung HS, O'Rourke AP, Agarwal SK. Nonoperative Management Is as Effective as Immediate Splenectomy for Adult Patients with High-Grade Blunt Splenic Injury. *J Am Coll Surg.* 2016 Aug;223(2):249-58.
6. Aoki M, Onogawa A, Matsumoto S, Matsushima K. Recent trends in the management of isolated high-grade splenic injuries: A nationwide analysis. *J Trauma Acute Care Surg.* 2023 Feb 1;94(2):220-225.
7. Schellenberg M, Ball CG, Owattanapanich N, Emigh B, Murphy PB, Moffat B, Mador B, Beckett A, Lee J, Joos E, Minor S, Strickland M, Inaba K; TAC Bile Leaks Study Group. Diagnosis and management of bile leaks after severe liver injury: A Trauma Association of Canada multicenter study. *J Trauma Acute Care Surg.* 2022 Dec 1;93(6):813-820.
8. Radding S, Harfouche MN, Dhillon NK, Ko A, Hawley KL, Kundi R, Maddox JS, Radowsky JS, DuBose JJ, Feliciano DV, Kozar RA, Scalea TM. A pseudo-dilemma: are we over-diagnosing and over-treating traumatic splenic intraparenchymal pseudoaneurysms? *J Trauma Acute Care Surg.* 2023 Aug 21.