

[Chronic Wasting Disease a Threat to Missouri Deer](#) April 22, 2014

nwf.org

... In 2010, captive deer at big-game hunting facilities in two Missouri counties were the first diagnosed with the fatal disease. Since then, wild deer near these facilities have also been diagnosed with the disease. ... special interests in Missouri are heavily pressuring for bills that would undermine the Missouri Department of Conservation's efforts [to manage the disease] —and possibly facilitate the spread of the disease. ...

[SCOMO to Hear Regulation Case as State Reports Increase in Deer Disease](#) January

9, 2018 OzarksFirst.com

... The state's Conservation Commission first detected chronic wasting disease at a private hunting preserve in 2010... it enacted regulations for importing and handling of the animals... A circuit court in mid Missouri's Gasconade County determined in 2016 that the commission lacked the authority to enforce the regulations. The Missouri Supreme Court will hear arguments in an appeal ...

[Chronic wasting disease gains more national attention](#) March 23, 2019 Missouri,

Lakenewsonline.com

... In Missouri ... the disease was first discovered in a captive deer facility in 2010 ... CWD has progressed slowly here, thanks in large part to the MDC aggressively culling in small, concentrated areas around the discovery sites... Wisconsin ... decided to basically do nothing about the disease and let it spread. They now have CWD in 56 counties with areas where 50 percent of bucks harvested are infected ...

These articles have been reviewed for relevance and content. Some links to the original article may no longer be active.