

# Tennis Elbow (Lateral Epicondylitis)

Tennis elbow, or *lateral epicondylitis*, is a painful condition of the elbow caused by overuse. Not surprisingly, playing tennis or other racquet sports can cause this condition. But several other sports and activities can also put you at risk.

Tennis elbow is a degeneration of the tendons that join the forearm muscles on the outside of the elbow. The forearm muscles and tendons become damaged from overuse — repeating the same motions again and again. This leads to pain and tenderness on the outside of the elbow.

There are many treatment options for tennis elbow. In most cases, treatment involves a team approach. Primary doctors, physical therapists, and, in some cases, surgeons work together to provide the most effective care.

## Anatomy

Your elbow joint is a joint made up of three bones: your upper arm bone (humerus) and the two bones in your forearm (radius and ulna). There are bony bumps at the bottom of the humerus called epicondyles. The bony bump on the outside (lateral side) of the elbow is called the lateral epicondyle. Muscles, ligaments, and tendons hold the elbow joint together.

Lateral epicondylitis, or tennis elbow, involves the muscles and tendons of your forearm. Your forearm muscles extend your wrist and fingers. Your forearm tendons -- often called extensors -- attach the muscles to bone. They attach on the lateral epicondyle. The tendon usually involved in tennis elbow is called the Extensor Carpi Radialis Brevis (ECRB).



Modified from *The Body Almanac*. © American Academy of Orthopaedic Surgeons, 2003.

## Cause

*Parts of handout from the AAOS. AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice.*

## Overuse

Recent studies show that tennis elbow is often due to damage to a specific forearm muscle. The extensor carpi radialis brevis (ECRB) muscle helps stabilize the wrist when the elbow is straight. This occurs during a tennis groundstroke, for example. When the ECRB is weakened from overuse, microscopic tears form in the tendon where it attaches to the lateral epicondyle. This leads to inflammation and pain.

The ECRB may also be at increased risk for damage because of its position. As the elbow bends and straightens, the muscle rubs against bony bumps. This can cause gradual wear and tear of the muscle over time.

## Activities

Athletes are not the only people who get tennis elbow. Many people with tennis elbow participate in work or recreational activities that require repetitive and vigorous use of the forearm muscle.

Painters, plumbers, and carpenters are particularly prone to developing tennis elbow. Studies have shown that auto workers, cooks, and even butchers get tennis elbow more often than the rest of the population. It is thought that the repetition and weight lifting required in these occupations leads to injury.

## Age

Most people who get tennis elbow are between the ages of 30 and 50, although anyone can get tennis elbow if they have the risk factors. In racquet sports like tennis, improper stroke technique and improper equipment may be risk factors.

## Unknown

Lateral epicondylitis can occur without any recognized repetitive injury. This occurrence is called "insidious" or of an unknown cause.

## Symptoms



Location of pain in lateral epicondylitis.

*Reproduced with permission from Griffen L (ed): Essentials of Musculoskeletal Care, Third Edition. © American Academy of Orthopaedic Surgeons, 2005.*

The symptoms of tennis elbow develop gradually. In most cases, the pain begins as mild and slowly worsens over weeks and months. There is usually no specific injury associated with the start of symptoms.

*Parts of handout from the AAOS. AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice.*

Common signs and symptoms of tennis elbow include:

- Pain or burning on the outer part of your elbow
- Weak grip strength

The symptoms are often worsened with forearm activity, such as holding a racquet, turning a wrench, or shaking hands. Your dominant arm is most often affected; however both arms can be affected.

## Doctor Examination

Your doctor will consider many factors in making a diagnosis. These include how your symptoms developed, any occupational risk factors, and recreational sports participation.

Your doctor will talk to you about what activities cause symptoms and where on your arm the symptoms occur. Be sure to tell your doctor if you have ever injured your elbow. If you have a history of rheumatoid arthritis or nerve disease, tell your doctor.

During the examination, your doctor will use a variety of tests to pinpoint the diagnosis. For example, your doctor may ask you to try to straighten your wrist and fingers against resistance with your arm fully straight to see if this causes pain. If the tests are positive, it tells your doctor that those muscles may not be healthy.

## Tests

Your doctor may recommend additional tests to rule out other causes of your problem.

### *X-rays*

These may be taken to rule out arthritis of the elbow.

### *Magnetic Resonance Imaging (MRI)*

If your doctor thinks your symptoms are related to a neck problem, an MRI scan may be ordered. This will help your doctor see if you have a possible herniated disk or arthritis in your neck. Both of these conditions often produce arm pain.

### *Electromyography (EMG)*

Your doctor may order an EMG to rule out nerve compression. Many nerves travel around the elbow, and the symptoms of nerve compression are similar to those of tennis elbow.

## Treatment

### *Nonsurgical Treatment*

Approximately 80% to 95% of patients have success with nonsurgical treatment.

**Rest.** The first step toward recovery is to give your arm proper rest. This means that you will have to stop participation in sports or heavy work activities for several weeks. All lifting should be done with **the palms up**.

**Non-steroidal anti-inflammatory medicines.** Drugs like aspirin or ibuprofen reduce pain and swelling.



Wrist stretching exercise with elbow extended.

**Equipment check.** If you participate in a racquet sport, your doctor may encourage you to have your equipment checked for proper fit. Stiffer racquets and looser-strung racquets often can reduce the stress on the forearm, which means that the forearm muscles do not have to work as hard. If you use an oversized racquet, changing to a smaller head may help prevent symptoms from recurring.

**Physical/Occupational therapy.** Specific exercises are helpful for strengthening the muscles of the forearm. Your therapist may also perform ultrasound, ice massage, or muscle-stimulating techniques to improve muscle healing.

**Brace.** Using a brace centered over the back of your forearm may also help relieve symptoms of tennis elbow. This can reduce symptoms by resting the muscles and tendons.



Counterforce brace.

### ***Surgical Treatment***

If your symptoms do not respond after 6 to 12 months of nonsurgical treatments, your doctor may recommend surgery.

Most surgical procedures for tennis elbow involve removing diseased muscle and reattaching healthy muscle back to bone.

The right surgical approach for you will depend on a range of factors. These include the scope of your injury, your general health, and your personal needs. Talk with your doctor about the options. Discuss the results your doctor has had, and any risks associated with each procedure.

**Open surgery.** The most common approach to tennis elbow repair is open surgery. This involves making an incision over the elbow.

*Parts of handout from the AAOS. AAOS does not endorse any treatments, procedures, products, or physicians referenced herein. This information is provided as an educational service and is not intended to serve as medical advice.*

Open surgery is usually performed as an outpatient surgery. It rarely requires an overnight stay at the hospital.

Arthroscopic surgery. Tennis elbow can also be repaired using tiny instruments and small incisions. Arthroscopic surgery is minimally invasive and may allow a faster recovery. Like open surgery, this is a same-day or outpatient procedure.

Surgical risks. As with any surgery, there are risks with tennis elbow surgery. The most common things to consider include:

- Infection
- Nerve and blood vessel damage
- Possible prolonged rehabilitation
- Loss of strength
- Loss of flexibility
- The need for further surgery

Rehabilitation. Following surgery, your arm may be immobilized temporarily with a splint. About 1 week later, the sutures and splint are removed.

After the splint is removed, exercises are started to stretch the elbow and restore flexibility. Light, gradual strengthening exercises are started about 2 months after surgery.

Your doctor will tell you when you can return to athletic activity. This is usually 4 to 6 months after surgery. Tennis elbow surgery is considered successful in 80% to 90% of patients. However, it is not uncommon to see a loss of strength.

For more information, please email me at [apastor@pssurgicalcenter.com](mailto:apastor@pssurgicalcenter.com).

Best wishes,

A handwritten signature in dark ink, appearing to read 'Andrew Pastor', with a stylized, flowing script.

Andrew Pastor, M.D.