

Unit 03: Differentiation Rules

Unit Objectives

- Learn the shortcuts for determining derivatives
 - Purchase coffin
 - Insert $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 - Bury deep in an abandoned field
- Find higher order derivatives
- Find derivatives of inverse functions

Unit 03 Lesson 01: Differentiate Polynomials

- Determine derivatives of polynomials using the Power Rule

~~~~~  
Can you spot the pattern?

~~~~~  
1. $\frac{d}{dx} 5x^3 =$

2. $\frac{d}{dx} 3x =$

3. $\frac{d}{dx} 3 =$

4. $\frac{d}{dx} \frac{2}{x} =$

5. $\frac{d}{dx} (2x^2 - \frac{5}{x^3}) =$

6. $\frac{d}{dx} (6\sqrt{x}) =$

~~~U03L01 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Power rule (positive integer powers)
2. [Khan] Differentiation: definition and basic derivative rules: Power rule (negative and fractional powers)

~~~U03L01 Classwork~~~

7. $\frac{d}{dx} \left(\frac{1}{6}x^2 - x \right) =$

8. $\frac{d}{dx} 3x^4 =$

9. $\frac{d}{dx} x =$

10. $\frac{d}{dx} 0 =$

11. $\frac{d}{dx} \frac{5}{x^2} =$

12. $\frac{d}{dx} \left(-\frac{5}{x} + 2x^2 \right) =$

13. $\frac{d}{dx} \frac{1}{6\sqrt{x}} =$



14. The height of a ball dropped from a 1000ft tall building can be found using the $y = -16t^2 + 1000$.
- Find the instantaneous velocity of the ball when it has been falling for 3 second.

- Find the instantaneous velocity of the ball when it has been falling for 4 seconds.

Unit 03 Lesson 02: Identify Compositions

Lesson Objectives

- Find a way to express a function as a composition of two functions

Identify $f(x)$ and $g(x)$ such that

- $h(x) = f(g(x))$ and
- $f(x)$ and $g(x)$ are functions you know how to differentiate.

Then determine $f'(x)$ and $g'(x)$ just for practice.

1. $h(x) = (4 - 3x)^3$

2. $h(x) = (x^2 - 2x)^2$

3. $h(x) = \sqrt{2x - 1}$

4. $h(x) = \left| \frac{x}{3} \right|$

5. $h(x) = \frac{2}{x - 1}$

6. $h(x) = (x+2)^3 - (x+2)^2 + 2(x+2) - 5$

~~~U03L02 Homework~~~

- [Khan] Differentiation: definition and basic derivative rules: Power rule (with rewriting the expression)
- [Khan] Differentiation: composite, implicit, and inverse functions: Identify composite functions

~~~U03L02 Classwork~~~

Identify  $f(x)$  and  $g(x)$  such that  $h(x) = f(g(x))$ .

1.  $h(x) = \sqrt{2x - 1}$

2.  $h(x) = \frac{1}{x^2 - x}$

3.  $h(x) = (-4 + \sin(x))^2$

4.  $h(x) = \ln\left(\frac{3}{x}\right)$

5.  $(x-3)^2 - (x-3) + \sqrt{x-3}$

6.  $h(x) = |x^4 - x|$

## Unit 03 Lesson 03: Differentiate Compositions (Chain Rule)

### Lesson Objectives

- Use the chain rule to differentiate composite functions

Chain Rule:

$$\frac{dy}{dx}$$

Determine  $\frac{dy}{dx}$ .

1.  $y = (4 - 3x)^3$

2.  $y = \sqrt{4 - x^3}$

3.  $y = \frac{4}{3x^2 + 1}$

4.  $y = (x+2)^3 - (x+2)^2 + 2(x+2) - 5$

### ~~~U03L03 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Basic derivative rules: find the error
2. [Khan] Differentiation: definition and basic derivative rules: Basic derivative rules: table

~~~U03L03 Classwork~~~

Determine y'

7. $y = \sqrt{2x - 1}$

8. $y = (x^2 - 2x)^2$

9. $y = \frac{4}{x^2 - x}$

10. $(5x-3)^2 - (4x-3) + \sqrt{2x - 3}$

11. $y = (5(2x^2-1)^2)^4$

12. $y = (4x^3 - 5(2x^2-1))^4$

Unit 03 Lesson 04: Differentiate Exponential Functions

Lesson Objectives

- Apply the differentiation rule for exponential functions

Differentiation Rule for exponential functions

Differentiate the following functions

1. $h(x) = 3(2^x) + 1$

2. $f(x) = 3e^x + 2x + 10$

3. $g(x) = e^{3x} + e^{x/4}$

4. $k(x) = e^{x^2 - x} - 2x + 5$

5. $m(x) = (x + e^{3x})^2$

~~~U03L04 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Differentiate polynomials

~~~U03L04 Classwork~~~

Differentiate the following functions

1.  $f(x) = 4x - 3e^x$

2.  $g(x) = 10^x + 5x + 4$

3.  $h(x) = 3x^4 - x^2 + e^{-2x}$

4.  $r(x) = e^{e^x}$

5.  $q(x) = \sqrt{e^{-3x} + 2}$



## Unit 03 Lesson 05: Differentiate Logarithmic Functions

### Lesson Objectives

- Apply the differentiation rule for exponential functions

### Differentiation Rule for exponential functions

Differentiate the following functions

1.  $h(x) = 2\log_{10}x$

2.  $f(x) = \ln(x+3) - 3x + 10$

3.  $g(x) = \ln(4x^2) + \ln(2)$

4.  $k(x) = \ln(3e^x) - 2x$

5.  $m(x) = (\ln(3x))^2$

### ~~~U03L05 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Differentiate integer powers (mixed positive and negative)

~~~U03L05 Classwork~~~

Differentiate the following functions

1. $f(x) = x^2 - x + \sqrt{x} - \ln(x) - e^x + 9$

2. $g(x) = -2\ln\left(\frac{1}{x^2}\right)$

3. $h(x) = \sqrt{\ln(x)} - 2x + 4$

4. $k(x) = 2\ln(2x + 4)$

5. Hmm, if $f'(x) = \frac{1}{2x + 5}$, what's the equation for $f(x)$?

Unit 03 Lesson 06: Differentiate Trigonometric Functions

Lesson Objectives

- Apply the differentiation rule for sine and cosine functions

Differentiation Rule for sine, cosine, and tangent functions

Differentiate the following equations

1. $y = 2\sin(x) + \tan(x) - 4$

2. $f(x) = 3\cos(2x) + 2x$

3. $g(x) = -\sin(x^2+1) + \ln(x) + e^x$

4. $z = \sin^2 x$

5. $y = \cos^3 x^2$

~~~U03L06 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Derivatives of $\sin(x)$ and $\cos(x)$
2. [Khan] Differentiation: definition and basic derivative rules: Derivatives of e^x and $\ln(x)$

~~~U03L06 Classwork~~~

Differentiate the following functions

1.  $f(x) = \tan(x) - 2x^3 + 1$

2.  $y = \cos^2 x$

3.  $g(x) = -\sin(2x) - \cos(3x) - x$

4.  $h(x) = \sin(\ln(x)) + \ln(x) + 10$

5.  $e^{\sin(x)} + 3x^2 - 5$

## Unit 03 Lesson 07: Differentiate Products of Functions

### Lesson Objectives

- Apply the differentiation rule for two functions multiplied together

### Product Rule

Differentiate the following equations

6.  $y = x^2 \sin x$

7.  $\ln(x) \cos(x) + 2x - e$

8.  $g(x) = e^x(x^2 - 2x)^3 - \pi x$

9.  $z = \sin x \cos 2x$

10.  $y = e^{2x} e^{-3x}$

### ~~~U03L07 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Differentiate products
2. [Khan] Differentiation: definition and basic derivative rules: Product rule with tables

~~~U03L07 Classwork~~~

Differentiate the following functions

6. $f(x) = 2x\cos x + 49$

7. $y = e^x \sin x + ex$

8. $g(x) = \cos^2 x$, solve two ways, a) rewrite as $(\cos x)^2$ b) rewrite as $\cos x \cos x$

9. $e^{-2x} \sin(-3x) - \pi x^2$

10. $e^{x \sin(x)} + 7$

Unit 03 Lesson 08: Differentiate Quotients of Functions

Lesson Objectives

- Apply the differentiation rule for two functions multiplied together

Quotient Rule

Differentiate the following equations

1. $y = \frac{x^2}{\sin x}$

2. $y = \frac{\cos x}{\ln x} + 4x - 2$

3. $f(x) = \frac{e^x}{(x^2 - 2x)^2} + e$

4. $f(x) = \frac{\cos 2x}{\sin x} + 54321$

5. $g(x) = \frac{x^3}{x^2}$, solve by simplifying first, then solve again without simplifying first

~~~U03L08 Homework~~~

1. [Khan] Differentiation: definition and basic derivative rules: Differentiate quotients
2. [Khan] Differentiation: definition and basic derivative rules: Quotient rule with tables

~~~U03L08 Classwork~~~

Differentiate the following functions

1.  $h(x) = \frac{\cos x}{2x}$

2.  $h(x) = \frac{\sin x}{e^x} + e^3$

3.  $h(x) = \frac{\sin x}{\cos x}$  solve by simplifying first, then solve again without simplifying first

4.  $y = \frac{e^{2x}}{\sin x}$

5.  $y = e^{\frac{\sin x}{x}}$




## Unit 03 Lesson 09: Differentiate Implicitly

### Lesson Objectives

- Differentiate equations that aren't solved for y.

1. A circle with radius 4 and center at (2, -1) has an equation of  $(x - 2)^2 + (y + 1)^2 = 16$ .

- a. Determine the slope of the tangent line at (-0.828, 1.828), and check by graphing.



- b. Determine the slope of the tangent line at (5.578, 0.789), and check by graphing.

- c. At what points will the tangent lines be horizontal?

- d. At what points will the tangent lines be vertical?

2. Find  $dy/dx$  of the equation  $x^3 + y^3 = 18xy$  and determine the slope of the tangent line at (2, 1)

Ans.  $dy/dx = (-2/11)$

3. Find  $dy/dx$  of the equation  $y^2 = \frac{x-1}{x+1}$  and determine the slope of the tangent line at (3, 0.25)  
Ans.  $dy/dx = 0.25$

4. Find  $dy/dx$  of the equation  $x = \sin y$  and determine the slope of the tangent line at (1,  $\pi/3$ )

Ans.  $dy/dx = 2$

5. Find  $dy/dx$  of the equation  $x + \sin y = xy$  and determine the slope of the tangent line at (2, 0)


Ans.  $dy/dx = 1$

~~~U03L09 Homework~~~

1. [Khan] Differentiation: composite, implicit, and inverse functions: Implicit differentiation
2. [Khan] Differentiation: definition and basic derivative rules: Differentiate rational functions

~~~U03L09 Classwork~~~

1. A circle with radius 5 and center at  $(-3, 2)$  has an equation of  $(x + 3)^2 + (y - 2)^2 = 25$ .
  - a. Determine the slope of the tangent line at  $(-6.536, -1.536)$ , and check by graphing.



- b. Determine the slope of the tangent line at  $(-5.236, 6.472)$ , and check by graphing.
  - c. At what points will the tangent lines be horizontal?
  - d. At what points will the tangent lines be vertical?

2. Find  $dy/dx$  of the equation  $x^2 + xy - y^2 = 3$  and determine the slope of the tangent line at (1, 1)

Ans.  $dy/dx = 3$

3. Find  $dy/dx$  of the equation  $x^2 = \frac{x - y}{x + y}$  and determine the slope of the tangent line at (3, 2)  
Ans.  $dy/dx = -24.33$

4. Find  $dy/dx$  of the equation  $x = \cos y$  and determine the slope of the tangent line at (1,  $\pi/6$ )

Ans.  $dy/dx = -2$

5. Find  $dy/dx$  of the equation  $x^{2/3} - y^{2/3} = 1$  and determine the slope of the tangent line at (64, 27)

Ans.  $dy/dx = 0.75$

## Unit 03 Lesson 10: Determine Higher Order Derivatives

### Lesson Objectives

- Differentiate more than once

### Notations:

- The rate of change of position is \_\_\_\_\_
- The rate of change of velocity is \_\_\_\_\_
- Therefore, the rate of change of the rate of change of position is \_\_\_\_\_



1. The height of a ball dropped from a 1000ft tall building can be found using the equation  $y = (-16\text{ft/sec}^2)t^2 + 1000\text{ft}$ .
  - a. Find the velocity of the ball when it has been falling for 2 seconds.

b. Find the velocity of the ball when it has been falling for 0 seconds.

c. Find the acceleration of the ball when it has been falling for 2 seconds.

d. Find the acceleration of the ball when it has been falling for 0 seconds.

2. A weight at  $y = 1\text{ft}$  is hanging peacefully on a spring, minding its own business. Alas, some wretched scoundrel comes and stretches it down 1 foot below the resting position, dooming the weight to an eternity of oscillation. At time  $t = 0\text{sec}$ , the weight is released. The height of the weight could be given by the equation  $f(t) = (-1\text{ft})\cos t + 1\text{ft}$

- a. What is the position of the weight at
  - i.  $t = 0$  seconds?
  - ii.  $t = \pi$  seconds?
  - iii. When is the first time the spring will return to its resting position?
  - iv. When is the first time the spring will return to the position it was released from?

- b. What is the velocity of the weight at
  - i.  $t = 0$  seconds?

- ii.  $t = \pi/2$  seconds?

- iii.  $t = \pi$  seconds?

- iv.  $t = 3\pi/2$  seconds?

- v.  $t = 2\pi$  seconds?


- c. What is the acceleration of the weight at
  - i.  $t = 0$  seconds?

- ii.  $t = \pi/2$  seconds?

- iii.  $t = \pi$  seconds?

- iv.  $t = 3\pi/2$  seconds?

- v.  $t = 2\pi$  seconds?



~~~U03L10 Homework~~~

1. [Khan] Differentiation: composite, implicit, and inverse functions: Second derivatives
2. [Khan] Differentiation: composite, implicit, and inverse functions: Second derivatives (implicit equations)

~~~U03L10 Classwork~~~

$$\frac{d^2 y}{dx^2}$$

1. Find  $\frac{d^2 y}{dx^2}$  if  $y = x \ln x$

2. Find  $f''(t)$  if  $f(t) = e^t \cos t$

3. The height of a ball dropped from a 300m tall building can be found using the equation  $y = (-9.81 \text{m/sec}^2)t^2 + 300\text{m}$ .

a. Find the velocity of the ball when it has been falling for 2 seconds.


b. Find the velocity of the ball when it has been falling for 0 seconds.

c. Find the acceleration of the ball when it has been falling for 2 seconds.

d. Find the acceleration of the ball when it has been falling for 0 seconds.

4. A weight at  $y = 0\text{m}$  is hanging peacefully on a spring, minding its own business. Alas, some wretched scoundrel comes and stretches it down 1 meter below the resting position, dooming the weight to an eternity of oscillation. At time  $t = 0\text{sec}$ , the weight is released. The height of the weight could be given by the equation  $f(t) = (-1\text{m})\cos(2\pi t)$

- a. What is the position of the weight at
  - i.  $t = 0$  seconds?
  - ii.  $t = 0.5$  seconds?
  - iii. When is the first time the spring will return to its resting position?
  - iv. When is the first time the spring will return to the position it was released from?
- b. What is the velocity of the weight at
  - i.  $t = 0$  seconds?
  - ii.  $t = 0.25$  seconds?
  - iii.  $t = 0.5$  seconds?
  - iv.  $t = 0.75$  seconds?
  - v.  $t = 1$  seconds?
- c. What is the acceleration of the weight at
  - i.  $t = 0$  seconds?
  - ii.  $t = 0.25$  seconds?
  - iii.  $t = 0.5$  seconds?
  - iv.  $t = 0.75$  seconds?
  - v.  $t = 1$  seconds?





## Unit 03 Lesson 11: Review for Differentiation Rules Quiz

### Lesson Objectives

- Prepare to get a grade on the quiz that would make Mama proud.

1. Differentiate the following functions

a.  $f(x) = 4\sqrt{x}$

b.  $f(x) = x \sin x$

c.  $f(x) = \cos^3 x$

d.  $f(x) = \frac{\ln x}{e^x}$

e.  $f(x) = 3\sqrt{x^2 + 4}$

2. For the equation  $y^2 + xy = x^2$  determine

a.  $\frac{dy}{dx}$


b.  $\frac{dx}{dy}$

3. For the equation  $y = \frac{\ln x}{x}$  determine  $y''$ . Simplify.

4. At  $t = 0$ , some guy kicks a block that happened to be sitting on an oiled surface, sending it sliding at 40ft/sec in the positive  $x$  direction. Ah, good ol' Blocky, friend of physics teachers around the world. Anyways, Blocky slides along the oiled surface with equation  $x = 40\ln(t+1)$  where  $x$  represents the horizontal distance traveled in feet, and  $t$  represents time in seconds. Answer the following questions.

\*extra credit: explain why each answer does or does not make physical sense

- a. What is the distance traveled when  $t = 0$ ?



- b. What is the distance traveled when  $t = 3$ ?

- c. What is the velocity of the block when  $t = 0$ ?

- d. What is the velocity of the block when  $t = 3$ ?

- e. What is the acceleration of the block when  $t = 0$ ?

- f. What is the acceleration of the block when  $t = 3$ ?