

HC Verma Solutions Rotational Mechanics

<http://www.learncbse.in/rotational-mechanics-hc-verma-concepts-physics-solutions/>
<https://goo.gl/3XPBG2>
<http://cbsetuts.blogspot.in/2018/01/hc-verma-solutions-rotational-mechanics.html>
<https://goo.gl/h39D3R>
<https://goo.gl/EzyLqs>
<https://goo.gl/s4QEF7>

Rotational Mechanics HC Verma Concepts of Physics Solutions

Rotational Mechanics HC Verma Concepts of Physics Solutions Chapter 10

- $\omega_0 = 0$; $\rho = 100 \text{ rev/s}$; $\omega = 2\pi$; $\rho = 200 \pi \text{ rad/s}$
 $\Rightarrow \omega = \omega_0 + \alpha t$
 $\Rightarrow \omega = \alpha t$
 $\Rightarrow \alpha = (200 \pi)/4 = 50 \pi \text{ rad/s}^2 \text{ or } 25 \text{ rev/s}^2$
 $\therefore \theta = \omega_0 t + 1/2 \alpha t^2 = 8 \times 50 \pi = 400 \pi \text{ rad}$
 $\therefore \alpha = 50 \pi \text{ rad/s}^2 \text{ or } 25 \text{ rev/s}^2$
 $\theta = 400 \pi \text{ rad.}$
- $\theta = 100 \pi$; $t = 5 \text{ sec}$
 $\theta = 1/2 \alpha t^2 \Rightarrow 100\pi = 1/2 \alpha \times 25$
 $\Rightarrow \alpha = 8\pi \times 5 = 40 \pi \text{ rad/s} = 20 \text{ rev/s}$
 $\therefore \alpha = 8\pi \text{ rad/s}^2 = 4 \text{ rev/s}^2$
 $\omega = 40\pi \text{ rad/s}^2 = 20 \text{ rev/s}^2.$
- Area under the curve will decide the total angle rotated


$\therefore$ maximum angular velocity = $4 \times 10 = 40 \text{ rad/s}$

Therefore, area under the curve = $1/2 \times 10 \times 40 + 40 \times 10 + 1/2 \times 40 \times 10$
 $= 800 \text{ rad}$

$\therefore$ Total angle rotated = 800 rad.
- $\alpha = 1 \text{ rad/s}^2$, $\omega_0 = 5 \text{ rad/s}$; $\omega = 15 \text{ rad/s}$
 $\therefore \omega = \omega_0 + \alpha t$
 $\Rightarrow t = (\omega - \omega_0)/\alpha = (15 - 5)/1 = 10 \text{ sec}$

Also, $\theta = \omega_0 t + 1/2 \alpha t^2$
 $= 5 \times 10 + 1/2 \times 1 \times 100 = 100 \text{ rad.}$
- $\theta = 5 \text{ rev}$, $\alpha = 2 \text{ rev/s}^2$, $\omega_0 = 0$; $\omega = ?$
 $\omega^2 = (2 \alpha \theta)$
 $\Rightarrow \omega = \sqrt{2 \times 2 \times 5} = 2\sqrt{5} \text{ rev/s.}$

or $\theta = 10\pi \text{ rad}$, $\alpha = 4\pi \text{ rad/s}^2$, $\omega_0 = 0$, $\omega = ?$



$$\omega^2 = (2 \alpha \theta)$$

$$\Rightarrow \omega = \sqrt{2 \times 2 \times 5} = 2\sqrt{5} \text{ rev/s.}$$

$$\text{or } \theta = 10\pi \text{ rad, } \alpha = 4\pi \text{ rad/s}^2, \omega_0 = 0, \omega = ?$$

$$\omega = \sqrt{2\alpha\theta} = 2 \times 4\pi \times 10\pi$$

$$= 4\pi\sqrt{5} \text{ rad/s} = 2\sqrt{5} \text{ rev/s.}$$

6. A disc of radius = 10 cm = 0.1 m

Angular velocity = 20 rad/s

$$\therefore \text{Linear velocity on the rim} = \omega r = 20 \times 0.1 = 2 \text{ m/s}$$

$$\therefore \text{Linear velocity at the middle of radius} = \omega r/2 = 20 \times (0.1)/2 = 1 \text{ m/s.}$$

7. $t = 1 \text{ sec, } r = 1 \text{ cm} = 0.01 \text{ m}$

$$\alpha = 4 \text{ rad/s}^2$$

$$\text{Therefore } \omega = \alpha t = 4 \text{ rad/s}$$

Therefore radial acceleration,

$$A_n = \omega^2 r = 0.16 \text{ m/s}^2 = 16 \text{ cm/s}^2$$

$$\text{Therefore tangential acceleration, } a_t = \alpha r = 0.04 \text{ m/s}^2 = 4 \text{ cm/s}^2.$$

8. The Block is moving the rim of the pulley

The pulley is moving at a $\omega = 10 \text{ rad/s}$

Therefore the radius of the pulley = 20 cm

Therefore linear velocity on the rim = tangential velocity = $r\omega$

$$= 20 \times 10 = 200 \text{ cm/s} = 2 \text{ m/s.}$$



- HC Verma Solutions,
- HC Verma,
- Solution of HC Verma Concept of Physics,
- Concepts of Physics Part 1 - HC Verma Solutions,
- HC Verma Concepts of Physics Solutions part 1 and part 2,
- HC Verma Solutions For Physics,
- hc verma part 1 pdf,
- h.c verma solutions free download full book,
- hc verma part 2 pdf,
- hc verma objective solutions,
- hc verma short answer solutions pdf,
- Physics HC Verma 1 - Solutions,
- Chapter wise solutions to HC Verma's Concepts of Physics,
- HC Verma Solutions Both Parts,
- HC VERMA SOLUTIONS (CHAPTERWISE)

Read more about [HC Verma Solutions Rotational Mechanics](#)