

**Box 1. List of papers eligible for the statistical analysis (involving the FMA-UE outcome measure).**

Beckelhimer et al. (2011)	1
Bhagubai et al. (2021)	2
Bui et al. (2022)	3
Bürge et al. (2008)	4
Bustrén et al. (2017)	5
Caimmi et al. (2022)	6
Carda et al. (2017)	7
Catalogna et al. (2023)	8
Chang et al. (2006)	9
Chang et al. (2021)	10
Chin et al. (2021a)	11
Chin et al. 2021b)	12
Choi et al. (2023)	13
Cirstea and Levin (2000)	14
Conroy et al. (2011)	15
Conroy et al. (2019)	16
Delavaran et al. (2017)	17
Demartino et al. (2019)	18
El-Nashar et al. (2019)	19
Feng et al. (2015)	20
Finley et al. (2005)	21
Finley et al. (2012)	22
Ghaziani et al. (2020)	23
Grigoras et al. (2016)	24
Grimm et al. (2016)	25
Hirakawa et al. (2018)	26
Iwamoto et al. (2022)	27
Jayasinghe et al. (2020)	28
Jeong et al. (2023)	29
Jiang et al. (2023)	30
Lee et al. (2022)	31
Liu et al. (2008)	32
Llorens et al. (2021)	33
Maenza et al. (2020)	34
Marquez-Chin et al. (2017)	35
Martins et al. (2015)	36
Mitrović et al. 2023)	37
Pang et al. (2006)	38
Duarte Pereira et al. (2015)	39
Plantin et al. (2021)	40
Plantin et al. (2022)	41
Pohl et al. (2000)	42
Rand et al. (2014)	43
Salazar et al. (2020)	44
Sanchez et al. (2006)	45

Sattar et al. (2023)	46
Schick et al. (2022)	47
Sebastián-Romagosa et al. (2020)	48
Shimodozono et al. (2014)	49
Shin et al. (2017)	50
Smith et al. (2018)	51
Stoykov et al. (2009)	52
Straudi et al. (2020a)	53
Straudi et al. (2020b)	54
Takebayashi et al. (2022)	55
Tanaka et al. (2019)	56
Veloza and Woodbury (2011)	57
Zondervan et al. (2015)	58

## **References (papers eligible for the statistical analysis - involving the FMA-UE outcome measure).**

1. Beckelhimer SC, Dalton AE, Richter CA, Hermann V, Page SJ. Computer-based rhythm and timing training in severe, stroke-induced arm hemiparesis. *American Journal of Occupational Therapy*. 2011;65(1):96-100. doi:10.5014/ajot.2011.09158
2. Bhagubai MMC, Wolterink G, Schwarz A, Held JPO, Van Beijnum BJF, Veltink PH. Quantifying Pathological Synergies in the Upper Extremity of Stroke Subjects with the Use of Inertial Measurement Units: A Pilot Study. *IEEE J Transl Eng Health Med*. 2021;9. doi:10.1109/JTEHM.2020.3042931
3. Bui KD, Lyn B, Roland M, Wamsley CA, Mendonca R, Johnson MJ. The Impact of Cognitive Impairment on Robot-Based Upper-Limb Motor Assessment in Chronic Stroke. *Neurorehabil Neural Repair*. 2022;36(9):587-595. doi:10.1177/15459683221110892
4. Bürge E, Kupper D, Finckh A, Ryerson S, Schnider A, Leemann B. Neutral Functional Realignment Orthosis Prevents Hand Pain in Patients With Subacute Stroke: A Randomized Trial. *Arch Phys Med Rehabil*. 2008;89(10):1857-1862. doi:10.1016/j.apmr.2008.03.023
5. Bustrén EL, Sunnerhagen KS, Alt Murphy M. Movement Kinematics of the Ipsilesional Upper Extremity in Persons with Moderate or Mild Stroke. *Neurorehabil Neural Repair*. 2017;31(4):376-386. doi:10.1177/1545968316688798

Ierardi, E., van Wijck, F., Ali, M., Best, C., Coupar, F., 2025 "Defining severity level descriptors for post-stroke upper limb motor impairment and activity limitation: a systematic review" – Journal Details

6. Caimmi M, Giovanzana C, Gasperini G, Molteni F, Molinari Tosatti L. Robot Fully Assisted Upper-Limb Functional Movements Against Gravity to Drive Recovery in Chronic Stroke: A Pilot Study. *Front Neurol.* 2022;12. doi:10.3389/fneur.2021.782094
7. Carda S, Biasiucci A, Maesani A, et al. Electrically Assisted Movement Therapy in Chronic Stroke Patients With Severe Upper Limb Paresis: A Pilot, Single-Blind, Randomized Crossover Study. *Arch Phys Med Rehabil.* 2017;98(8):1628-1635.e2. doi:10.1016/j.apmr.2017.02.020
8. Catalogna M, Hadanny A, Parag Y, Adler M, Elkarif V, Efrati S. Functional MRI evaluation of hyperbaric oxygen therapy effect on hand motor recovery in a chronic post-stroke patient: a case report and physiological discussion. *Front Neurol.* 2023;14. doi:10.3389/fneur.2023.1233841
9. Chang JJ, Tung WL, Wu WL, Su FC. Effect of bilateral reaching on affected arm motor control in stroke - with and without loading on unaffected arm. *Disabil Rehabil.* 2006;28(24):1507-1516.  
<http://search.ebscohost.com/login.aspx?direct=true&db=amed&AN=0094605&site=ehost-live&scope=site>
10. Chang JL, Coggins AN, Saul M, et al. Transcutaneous Auricular Vagus Nerve Stimulation (tAVNS) Delivered During Upper Limb Interactive Robotic Training Demonstrates Novel Antagonist Control for Reaching Movements Following Stroke. *Front Neurosci.* 2021;15. doi:10.3389/fnins.2021.767302
11. Chin LF, Hayward KS, Chai ALM, Brauer SG. A Self-Empowered Upper Limb Repetitive Engagement Program to Improve Upper Limb Recovery Early

Ierardi, E., van Wijck, F., Ali, M., Best, C., Coupar, F., 2025 "Defining severity level descriptors for post-stroke upper limb motor impairment and activity limitation: a systematic review" – Journal Details

Post-Stroke: Phase II Pilot Randomized Controlled Trial. *Neurorehabil Neural Repair*. 2021;35(9):836-848. doi:10.1177/15459683211032967

12. Chin LF, Hayward KS, Brauer SG. Factors Influencing Paretic Upper Limb Use during First 4 Weeks after Stroke: A Cross-Sectional Accelerometry Study. *Am J Phys Med Rehabil*. 2021;100(2):153-160. doi:10.1097/PHM.0000000000001539
13. Choi H, Park D, Rha DW, Nam HS, Jo YJ, Kim DY. Kinematic analysis of movement patterns during a reach-and-grasp task in stroke patients. *Front Neurol*. 2023;14. doi:10.3389/fneur.2023.1225425
14. Cirstea MC, Levin MF. Compensatory strategies for reaching in stroke. *Brain*. 2000;123(Pt 5):940-953. doi:10.1093/brain/123.5.940
15. Conroy SS, Whitall J, Dipietro L, et al. Effect of gravity on robot-assisted motor training after chronic stroke: A randomized trial. *Arch Phys Med Rehabil*. 2011;92(11):1754-1761. doi:10.1016/j.apmr.2011.06.016
16. Conroy SS, Wittenberg GF, Krebs HI, Zhan M, Bever CT, Whitall J. Robot-Assisted Arm Training in Chronic Stroke: Addition of Transition-to-Task Practice. *Neurorehabil Neural Repair*. 2019;33(9):751-761. doi:10.1177/1545968319862558
17. Delavaran H, Aked J, Sjunnesson H, et al. Spontaneous Recovery of Upper Extremity Motor Impairment After Ischemic Stroke: Implications for Stem Cell-Based Therapeutic Approaches. *Transl Stroke Res*. 2017;8(4):351-361. doi:10.1007/s12975-017-0523-9

18. Demartino AM, Rodrigues LC, Gomes RP, Michaelsen SM. Hand function and type of grasp used by chronic stroke individuals in actual environment. *Top Stroke Rehabil.* 2019;26(4):247-254. doi:10.1080/10749357.2019.1591037
19. El-Nashar H, Elwishy A, Helmy H, El-Rwainy R. Do core stability exercises improve upper limb function in chronic stroke patients? *Egyptian Journal of Neurology, Psychiatry and Neurosurgery.* 2019;55(1):1-9. doi:10.1186/s41983-019-0087-6
20. Feng W, Wang J, Chhatbar PY, et al. Corticospinal tract lesion load: An imaging biomarker for stroke motor outcomes. *Ann Neurol.* 2015;78(6):860-870. doi:10.1002/ana.24510
21. Finley MA, Fasoli SE, Dipietro L, et al. Short-duration robotic therapy in stroke patients with severe upper-limb motor impairment. *J Rehabil Res Dev.* 2005;42(5):683-691. doi:10.1682/JRRD.2004.12.0153
22. Finley M, Combs S, Carnahan K, Peacock S, Buskirk A Van. Comparison of "less affected limb" reaching kinematics in individuals with chronic stroke and healthy age-matched controls. *Phys Occup Ther Geriatr.* 2012;30(3):245-259. doi:10.3109/02703181.2012.716506
23. Ghaziani E, Couppé C, Siersma V, et al. Easily Conducted Tests During the First Week Post-stroke Can Aid the Prediction of Arm Functioning at 6 Months. *Front Neurol.* 2020;10(1371):1-13. doi:10.3389/fneur.2019.01371
24. Grigoras AV, Irimia DC, Poboroniuc MS, Popescu CD. Testing of a hybrid FES-robot assisted hand motor training program in sub-acute stroke survivors. *Advances in*

*Electrical and Computer Engineering*. 2016;16(4):89-94.

doi:10.4316/AECE.2016.04014

25. Grimm F, Walter A, Spüler M, Naros G, Rosenstiel W, Gharabaghi A. Hybrid neuroprosthesis for the upper limb: Combining brain-controlled neuromuscular stimulation with a multi-joint arm exoskeleton. *Front Neurosci*. 2016;10(367):1-11. doi:10.3389/fnins.2016.00367
26. Hirakawa Y, Takeda K, Tanabe S, et al. Effect of intensive motor training with repetitive transcranial magnetic stimulation on upper limb motor function in chronic post-stroke patients with severe upper limb motor impairment. *Top Stroke Rehabil*. 2018;25(5):321-325. doi:10.1080/10749357.2018.1466971
27. Iwamoto Y, Imura T, Tanaka R, et al. Clinical Prediction Rule for Identifying the Stroke Patients who will Obtain Clinically Important Improvement of Upper Limb Motor Function by Robot-Assisted Upper Limb. *J Stroke Cerebrovasc Dis*. 2022;31(7):106517. doi:10.1016/j.jstrokecerebrovasdis.2022.106517
28. Jayasinghe SAL, Good D, Wagstaff DA, Winstein C, Sainburg RL. Motor Deficits in the Ipsilesional Arm of Severely Paretic Stroke Survivors Correlate With Functional Independence in Left, but Not Right Hemisphere Damage. *Front Hum Neurosci*. 2020;14. doi:10.3389/fnhum.2020.599220
29. Jeong EJ, Kang MJ, Lee S, et al. Predictors of manual dexterity at 3 and 6 months after stroke: Integration of clinical, neurophysiological, and neuroimaging factors. *International Journal of Rehabilitation Research*. 2023;46(4):308-315. doi:10.1097/MRR.0000000000000601

Ierardi, E., van Wijck, F., Ali, M., Best, C., Coupar, F., 2025 "Defining severity level descriptors for post-stroke upper limb motor impairment and activity limitation: a systematic review" – Journal Details

30. Jiang Y, Liu Z, Liu T, Ma M, Tang M, Chai Y. A Serious Game System for Upper Limb Motor Function Assessment of Hemiparetic Stroke Patients. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*. 2023;31:2640-2653.  
doi:10.1109/TNSRE.2023.3281408
31. Lee J, Kim H, Kim J, Chang WH, Kim YH. Multimodal Imaging Biomarker-Based Model Using Stratification Strategies for Predicting Upper Extremity Motor Recovery in Severe Stroke Patients. *Neurorehabil Neural Repair*. 2022;36(3):217-226.  
doi:10.1177/15459683211070278
32. Liu W, Mukherjee M, Sun C, Liu H, McPeak LK. Electroacupuncture may help motor recovery in chronic stroke survivors: A pilot study. *J Rehabil Res Dev*. 2008;45(4):587-596. doi:10.1682/JRRD.2007.11.0181
33. Llorens R, Fuentes MA, Borrego A, et al. Effectiveness of a combined transcranial direct current stimulation and virtual reality-based intervention on upper limb function in chronic individuals post-stroke with persistent severe hemiparesis: a randomized controlled trial. *J Neuroeng Rehabil*. 2021;18(1). doi:10.1186/s12984-021-00896-2
34. Maenza C, Good DC, Winstein CJ, Wagstaff DA, Sainburg RL. Functional Deficits in the Less-Impaired Arm of Stroke Survivors Depend on Hemisphere of Damage and Extent of Paretic Arm Impairment. *Neurorehabil Neural Repair*. 2020;34(1):39-50.  
doi:10.1177/1545968319875951
35. Marquez-Chin C, Bagher S, Zivanovic V, Popovic MR. Functional electrical stimulation therapy for severe hemiplegia: Randomized control trial revisited.

Ierardi, E., van Wijck, F., Ali, M., Best, C., Coupar, F., 2025 "Defining severity level descriptors for post-stroke upper limb motor impairment and activity limitation: a systematic review" – Journal Details

*Canadian Journal of Occupational Therapy*. 2017;84(2):87-97.

doi:10.1177/0008417416668370

36. Martins JC, Aguiar LT, Lara EM, Teixeira-Salmela LF, Faria CDC de M. Assessment of grip strength with the modified sphygmomanometer test: Association between upper limb global strength and motor function. *Braz J Phys Ther*. 2015;19(6):498-506. doi:10.1590/bjpt-rbf.2014.0118
37. Mitrović SZ, Konstantinović LM, Miler Jerković V, Dedijer-Dujović S, Djordjević OC. Extended Poststroke Rehabilitation Combined with Cerebrolysin Promotes Upper Limb Motor Recovery in Early Subacute Phase of Rehabilitation: A Randomized Clinical Study. *Medicina (Lithuania)*. 2023;59(2). doi:10.3390/medicina59020291
38. Pang MY, Harris JE, Eng JJ. A community-based upper-extremity group exercise program improves motor function and performance of functional activities in chronic stroke: A randomized controlled trial. *Arch Phys Med Rehabil*. 2006;87(1):1-9. doi:10.1016/j.apmr.2005.08.113
39. Duarte Pereira N, Vieira L, Pompeu FP, Menezes IDS, Anjos SM Dos, Ovando AC. Translation, cultural adaptation and reliability of the brazilian version of the Graded Wolf Motor Function Test in adults with severe hemiparesis. *Fisioterapia em Movimento*. 2015;28(4):667-676. doi:10.1590/0103-5150.028.004.ao03
40. Plantin J, Verneau M, Godbolt AK, et al. Recovery and Prediction of Bimanual Hand Use After Stroke. *Neurology*. 2021;97(7):E706-E719. doi:10.1212/WNL.0000000000012366

41. Plantin J, Godbolt AK, Pennati G V., et al. Motor inhibition and its contribution to recovery of dexterous hand use after stroke. *Brain Commun.* 2022;4(5).  
doi:10.1093/braincomms/fcac241
42. Pohl PS, Luchies CW, Stoker-Yates J, Duncan PW. Upper Extremity Control in Adults Post Stroke with Mild Residual Impairment. *Neurorehabil Neural Repair.* 2000;14(1):33-41.
43. Rand D, Givon N, Weingarden H, Nota A, Zeilig G. Eliciting upper extremity purposeful movements using video games: A comparison with traditional therapy for stroke rehabilitation. *Neurorehabil Neural Repair.* 2014;28(8):733-739.  
doi:10.1177/1545968314521008
44. Salazar AP, Cimolin V, Schifino GP, Rech KD, Marchese RR, Pagnussat AS. Bi-cephalic transcranial direct current stimulation combined with functional electrical stimulation for upper-limb stroke rehabilitation: A double-blind randomized controlled trial. *Ann Phys Rehabil Med.* 2020;63(1):4-11. doi:10.1016/j.rehab.2019.05.004
45. Sanchez RJ, Liu J, Rao S, et al. Automating arm movement training following severe stroke: Functional exercises with quantitative feedback in a gravity-reduced environment. *IEEE Transactions on Neural Systems and Rehabilitation Engineering.* 2006;14(3):378-389. doi:10.1109/TNSRE.2006.881553
46. Sattar E, Ahmad U, Rehman M, et al. Impaired Upper Limb Motor Function in Post-stroke Patients and Its Impact on Trunk Control and Mobility: A Cross-sectional Study. *Iranian Rehabilitation Journal.* 2023;21(2):283-290.  
doi:10.32598/irj.21.2.1834.1

47. Schick T, Kolm D, Leitner A, Schober S, Steinmetz M, Fheodoroff K. Efficacy of Four-Channel Functional Electrical Stimulation on Moderate Arm Paresis in Subacute Stroke Patients—Results from a Randomized Controlled Trial. *Healthcare*. 2022;10(4):704. doi:10.3390/healthcare10040704
48. Sebastián-Romagosa M, Cho W, Ortner R, et al. Brain Computer Interface Treatment for Motor Rehabilitation of Upper Extremity of Stroke Patients—A Feasibility Study. *Front Neurosci*. 2020;14. doi:10.3389/fnins.2020.591435
49. Shimodozono M, Noma T, Matsumoto S, Miyata R, Etoh S, Kawahira K. Repetitive facilitative exercise under continuous electrical stimulation for severe arm impairment after sub-acute stroke: A randomized controlled pilot study. *Brain Inj*. 2014;28(2):203-210. doi:10.3109/02699052.2013.860472
50. Shin JH, Park G, Cho DY. Cognitive-Motor Interference on Upper Extremity Motor Performance in a Robot-Assisted Planar Reaching Task Among Patients With Stroke. *Arch Phys Med Rehabil*. 2017;98(4):730-737. doi:10.1016/j.apmr.2016.12.004
51. Smith C, Kenney L, Howard D, et al. Prediction of setup times for an advanced upper limb functional electrical stimulation system. *J Rehabil Assist Technol Eng*. 2018;5:1-9. doi:10.1177/2055668318802561
52. Stoykov ME, Lewis GN, Corcos DM. Comparison of bilateral and unilateral training for upper extremity hemiparesis in stroke. *Neurorehabil Neural Repair*. 2009;23(9):945-953. doi:10.1177/1545968309338190

53. Straudi S, Baroni A, Mele S, et al. Effects of a Robot-Assisted Arm Training Plus Hand Functional Electrical Stimulation on Recovery After Stroke: A Randomized Clinical Trial. *Arch Phys Med Rehabil.* 2020;101(2):309-316.  
doi:10.1016/j.apmr.2019.09.016
54. Straudi S, Baroni A, Mele S, et al. Effects of a Robot-Assisted Arm Training Plus Hand Functional Electrical Stimulation on Recovery After Stroke: A Randomized Clinical Trial. *Arch Phys Med Rehabil.* 2020;101(2):309-316.  
doi:10.1016/j.apmr.2019.09.016
55. Takebayashi T, Takahashi K, Amano S, et al. Robot-Assisted Training as Self-Training for Upper-Limb Hemiplegia in Chronic Stroke: A Randomized Controlled Trial. *Stroke.* 2022;53(7):2182-2191.  
doi:10.1161/STROKEAHA.121.037260
56. Tanaka T, Hamaguchi T, Suzuki M, et al. Estimation of Motor Impairment and Usage of Upper Extremities during Daily Living Activities in Poststroke Hemiparesis Patients by Observation of Time Required to Accomplish Hand Dexterity Tasks. *Biomed Res Int.* 2019;2019(9471921):1-8.
57. Velozo CA, Woodbury ML. Translating measurement findings into rehabilitation practice: An example using Fugl-Meyer Assessment-Upper Extremity with patients following stroke. *J Rehabil Res Dev.* 2011;48(10):1211-1222.  
doi:10.1682/JRRD.2010.10.0203
58. Zondervan DK, Augsburger R, Bodenhofer B, Friedman N, Reinkensmeyer DJ, Cramer SC. Machine-based, self-guided home therapy for individuals with severe

Ierardi, E., van Wijck, F., Ali, M., Best, C., Coupar, F., 2025 "Defining severity level descriptors for post-stroke upper limb motor impairment and activity limitation: a systematic review" – Journal Details

arm impairment after stroke: A randomized controlled trial. *Neurorehabil Neural*

*Repair*. 2015;29(5):395-406. doi:10.1177/1545968314550368