



SCOTTSDALE FAMILY

Dentistry & Orthodontics

Surgical Implant Informed Consent

Because of the wide differences among people and dental conditions, a successful outcome cannot always be obtained. Dentistry is not an exact science and no guarantees or assurance as to the outcome of treatment or surgery can be made. My dentist has examined my mouth and alternatives to this treatment have been explained. To my knowledge, I have given an accurate report of my physical/mental health history to my Dentist. I have also reported any prior allergic or unusual reactions, abnormal bleeding or any other conditions related to my health.

I have been informed of the possible risks and such complications include:

- Pain, swelling, infection and discoloration and non integration of the implant(s)
- Numbness of the lip, tongue, chin, cheek, teeth or other areas may occur (The exact duration may not be determinable and may be irreversible).
- Inflammation or injury to the adjacent teeth present, bone fractures, sinus penetration, delayed healing, and allergic reactions to drugs or medications used.

Excessive smoking, alcohol, sugar, systemic disease and certain medications may affect gum healing and limit the success of the implant. I agree to follow my doctor's home care instructions and will report to my doctor for regular examinations as instructed.

I understand what is necessary to accomplish the placement of the implant under the gum and in the bone. Recognizing there is no method to accurately predict the gum and/or bone healing capabilities following the placement of the implant, and it has been explained that in some instances implants fail and must be removed.

I request and authorize permission for implant surgery and fully understand that during and/or following the treatment, conditions may become apparent which warrant, in the judgment of the doctor, additional or alternative treatment pertinent to the success of the implant. I also give authorization to release patient information to other healthcare providers, insurance companies and other involved parties as deemed appropriate by your office. I also approve any modifications in the procedure or materials used if it is felt necessary.

Dental Surgery & Risks Associated with History of Bisphosphonate Use

If you are now or have previously been treated by your physician with a class of medication known as bisphosphonates, you may be at risk for developing bisphosphonate related osteonecrosis of the jaw (BRONJ) or medication related osteonecrosis of the jaw (MRONJ).

- **What is bisphosphonate related osteonecrosis of the jaw (BRONJ)?**
BRONJ is a condition in which the bone of the upper or lower jaw does not heal properly and can remain exposed to the oral cavity. The lack of healing with BRONJ is typically related to impaired blood supply or disruption of the cells that help bone heal.
- **What is the risk for developing BRONJ if I am taking bisphosphonate medications?**

Current research estimates that the risk of developing BRONJ varies depending on the type of bisphosphonate medication you are taking.

- If you are taking injectable or *intravenous bisphosphonate medications*, the risk of developing BRONJ in association with certain dental procedures can be as high as 28%.*
- If you are taking *oral bisphosphonate medications*, the risk of developing BRONJ in association with certain dental procedures is typically less than 0.1%.*

**These risks quoted in this consent form are based on current research, but are not guaranteed to be the most current estimates as numerous and new studies relating to BRONJ are frequently published.*

- **Which dental procedures have been associated with the development of BRONJ?**

Dental procedures involving exposure of bone or penetration of the gum tissue have been shown to have a high association with the development of BRONJ. The procedure most commonly associated with the development of BRONJ is the removal (extraction) of teeth. Any surgical procedure may increase your risk of developing BRONJ.

- **If I stop taking bisphosphonate medication for a short time prior to my dental procedure, will that reduce my risk of developing BRONJ?**

The half-life of bisphosphonate medications has been shown to be anywhere from 7 to 10 years. Depending on the type of bisphosphonate medication you are taking (oral, injectable, or intravenous), discontinuation of your bisphosphonate medication may not reduce your risk of developing BRONJ and may actually put you at higher risk for injury to other parts of your body. Changes to your bisphosphonate regimen should not be made without strict consultation with your treating physician.

- **Are there any tests to determine if I may be treated safely?**

Currently, the only test indicated in medical/dental literature to assist with assessing risk for the development of BRONJ is the serum C-terminal telopeptide (CTX) test. This is a simple blood test that measures markers in the blood that are associated with remodeling of bone. Please note that while CTX blood tests are one potential indicator of risk for BRONJ susceptibility, this test is not currently seen as a foolproof measure to predict BRONJ risk and a this condition may still result even results of this test are satisfactory.

- **What are the signs and symptoms of BRONJ?**

You should notify your dentist immediately if you have any of the following symptoms now or anytime following your treatment. In some cases, patients have developed BRONJ months or even years after their dental procedure.

Numbness of the jaw

Pain in the jaw

Swelling of the jaw

Loosening teeth

Drainage

Exposed bone

- **How can I reduce my risk of BRONJ?**

Recent updates provided by the American Academy of Oral and Maxillofacial Surgeons (Oct 2022) indicates that no particular strategy can completely eliminate MRONJ/BRONJ risk, but the following can reduce this risk: 1) quitting smoking; 2) having optimal diabetes control; 3) maintaining good oral hygiene.

I hereby certify that staff have presented this information to aid in my decision-making process. I have been given the opportunity to ask the doctor and/or staff questions I have about the proposed dental treatment and the risks associated with implants, bisphosphonate use and the development of BRONJ. I have fully disclosed my current or past use of bisphosphonate medications to the doctors and/or their staff. I understand the recommended treatment, the fee(s) involved, the risks of treatment, any alternatives and risks of those alternatives, including the consequences of doing nothing. I have had all of my questions answered, and have not been offered any guarantees.

Patient Name _____ **Tooth/Area** _____

Patient or Guardian Signature _____ **Date** _____

Witness Signature _____ **Date** _____