



STUDI ANTIFUNGI TERHADAP *Curvularia* sp. DARI EKSTRAK DAUN GAMBIR (*Uncaria gambir* Roxb.)

ANTIFUNGAL STUDY AGAINST *Curvularia* sp. FROM GAMBIR LEAF EXTRACT (*Uncaria gambir* Roxb.)

(Font times New Roman-14, Centered & Bold)

Rexi Iswanto ¹, Dodi Iskandar ^{1*}, Zaenal Mutaqin ¹, Muliani ¹, Danie Indra Yama ¹

¹ Nursing and caregiving support services, SMK Muhammadiyah 1 Kuburaya, Kuburaya

*email Koresponden: iskandar.dodi79@gmail.com

Abstract

This study aims to determine the effect of gambir leaf extract on the growth of *Curvularia* sp. fungus and to determine the concentration of extract that can inhibit 50% of the growth of fungal colonies. The research method used was a Completely Randomized Design (CRD) with 4 variations of extract concentration (0%, 5%, 10%, and 15%), each repeated 6 times. The results showed that the highest concentration of gambir leaf extract in inhibiting fungal growth was at a concentration of 15%. LC50% analysis revealed that the concentration required to achieve 50% inhibition was 118.16%, indicating that the effectiveness of the extract against *Curvularia* sp. fungus is low or less toxic.

Keywords: Gambir leaves, *Curvularia* sp., antifungal

Abstrak

Penelitian ini bertujuan untuk mengetahui pengaruh ekstrak daun gambir terhadap pertumbuhan jamur *Curvularia* sp. dan menentukan konsentrasi ekstrak yang mampu menghambat 50% pertumbuhan koloni jamur. Metode penelitian yang digunakan adalah Rancangan Acak Lengkap (RAL) dengan 4 variasi konsentrasi ekstrak (0%, 5%, 10%, dan 15%) yang masing-masing diulang sebanyak 6 kali. Hasil penelitian menunjukkan bahwa konsentrasi ekstrak daun gambir tertinggi dalam menghambat pertumbuhan jamur terdapat pada konsentrasi 15%. Analisis LC50% diketahui bahwa konsentrasi yang dibutuhkan untuk mencapai daya hambat 50% adalah 118,16%, hal ini menunjukkan bahwa efektivitas ekstrak terhadap jamur *Curvularia* sp. tergolong rendah atau kurang toksik

Kata Kunci : Daun gambir, *Curvularia* sp., antijamur

INTRODUCTION

Curvularia sp. is one type of pathogenic fungus that is often found in plantation crops. This fungus is known as a causative agent of various plant diseases, such as leaf spots, which can have a negative impact on plant productivity. Control of this fungus is generally carried out using



synthetic fungicides. However, long-term use of chemical fungicides can cause resistance in fungi and adverse side effects (Anugrah, & Widiyanti, 2018).

Along with increasing awareness of the importance of environmentally friendly and sustainable agriculture, the search for alternative biological control is becoming increasingly important. One potential alternative is to use natural compounds from plants that have antifungal activity. One plant that shows potential as an antifungal is *Uncaria gambir* Roxb. which is known as gambir.

Gambir leaves have long been used in various traditional medicines in Southeast Asia, mainly due to their bioactive compounds such as catechins. Previous studies have shown that gambir leaf extract has various biological activities, including antifungal activity such as *Candida albicans* (Rosa, 2021). However, specific studies on the effectiveness of gambir leaves as an antifungal against *Curvularia* sp. are still limited. This study aims to explore the potential of gambir leaf extract as an antifungal in inhibiting the growth of *Curvularia* sp. with various extract concentrations.

RESEARCH METHODS

The design used was a Completely Randomized Design (CRD) with four treatments of gambir extract concentration: 0% (negative control), 5%, 10%, and 15%, which were repeated six times, resulting in 24 experimental units.

Tools

The tools used in this study were autoclave, sieve, stirring rod, spray bottle, vial bottle, petri dish, porcelain dish, cork borer, Erlenmeyer flask, beaker, measuring cup, hammer mill, hotplate magnetic stirrer, incubator, vernier caliper, ose needle, refrigerator, laminar air flow (LAF), microscope, Bunsen burner, tweezers, dropper pipette, oven, rotary vacuum evaporator, spatula, test tube, digital scale, glass jar, vortex, and waterbath.

Materials

The materials used in this study were agar, aluminum foil, distilled water, 70% alcohol, acetic acid (CH₃COOH), sulfuric acid (H₂SO₄), gambir leaves (*Uncaria gambir* Roxb.), 96% ethanol, 10% FeCl₃, granulated sugar, 2 N HCl, 0.1 M dilute HCl, *Curvularia* sp. isolate, cotton, potatoes, filter paper, 10% NaOH, heat-resistant plastic, plastic wrap, ciulei reagent, mayer reagent, wagner reagent, spirits, and tissue.

Procedure

In Silico Test



Using the KNApSAcK website (<http://www.knapsackfamily.com>) and PASS Online (www.way2drug.com) to identify the antifungal potential of compounds in *Uncaria gambir* based on SMILES codes and biological activity probability.

Making Simplisia

Gambir leaves are dried using sunlight, ground into powder using a hammer mill, and stored in a container protected from direct sunlight.

Extract Preparation

Following the maceration method (Safitri, Putra, & Irawan, 2023), 500 grams of gambir leaf powder was soaked in 96% ethanol (1:4) for 72 hours, filtered, and concentrated using a rotary evaporator.

Ethanol-Free Test

To detect the presence of ethanol, 1 ml of extract was tested with acetic acid and sulfuric acid.

Phytochemical Test

Screening for active compounds such as alkaloids, flavonoids, tannins, terpenes, and saponins was carried out to ensure the content of bioactive compounds.

Sterilization of Tools and Materials

All tools and materials were sterilized by autoclaving at 121°C for 15 minutes or by Bunsen burner.

Preparation of PDA Media

Potato dextrose agar (PDA) media was made by boiling potatoes, mixing them with agar and sugar, then putting them in an Erlenmeyer flask and sterilizing them.

Isolate Rejuvenation

Isolate *Curvularia* sp. rejuvenated on PDA media before toxicity test to ensure active metabolism.

Toxicity Test

Using the food poisoned technique by adding gambir leaf extract (5%, 10%, and 15%) to PDA media and measuring the growth of *Curvularia* sp. fungus after incubation.

Data Analysis

The data collected are primary data obtained by measuring the diameter of mushroom growth using a caliper (mm). The data are then tabulated and analyzed using Analysis of Variance (ANOVA) with Minitab software. If there is a significant difference, further analysis will be carried out using Tukey's multiple comparison test at a 99% confidence level.



Sub-bab (optional, left navigation and bold)

RESULTS AND DISCUSSION

Analysis of Compound Potential through In Silico Test

The results of the in silico test show that most compounds in gambir leaves have moderate biological potential as antifungal agents, with an active probability (Pa) above 0.3. Compounds such as Cinchonain Ia, Gambiriin B1, and Catechin are among those that have potential for further research, while Mitraphylline and Uncarine B show low potential.

Gambir Leaf Extract Yield

Gambir leaf extraction produces a yield of 16%, exceeding the minimum standard of 10% based on the Herbal Pharmacopoeia. This high yield indicates the optimal active substance content for further research as an antifungal agent.

Ethanol-Free Test Results

The ethanol-free test showed that the gambir leaf extract did not contain ethanol, ensuring that the antifungal activity produced came entirely from the active compounds in the gambir leaves, not from ethanol as a solvent.

Phytochemical Test

The phytochemical test showed the presence of phenols, terpenoids, and saponins in the gambir leaf extract, which are known to have antimicrobial activity. The absence of alkaloids and flavonoids is likely due to the excessive drying process. The presence of phenols, terpenoids, and saponins strengthens the potential of gambir leaves as an antifungal agent.

Antifungal Test

At a concentration of 15%, the highest inhibition was achieved with an average of 34.93% on the 2nd day after incubation, which then decreased to 23.59% on the 8th day. This is thought to be due to the degradation of active compounds over time or the adaptation of fungi to the environment.

The results of statistical tests showed that gambir leaf extract provided significant inhibition of *Curvularia* sp., with a significant effect on increasing extract concentration. Antifungal compounds such as terpenoids, saponins, and phenols in the extract play a role in disrupting cell membranes and the fungal reproduction process, although their effectiveness is limited.

Probit analysis showed that the LC50% value of this extract reached 118.16%, indicating low toxicity and the need for high concentrations to inhibit 50% of fungal growth. The low effectiveness of the extract was due to the minimal number of active compounds such as alkaloids and flavonoids detected in the extract, indicating that the extraction process or preparation method may need to be refined to increase the bioactive content.

CONCLUSION

This study examined the antifungal effects of gambir leaf extract on *Curvularia* sp. using the food poisoning method. The results showed that gambir leaf extract was able to inhibit the



growth of *Curvularia* sp., with the highest inhibition at a concentration of 15%, although its overall effectiveness was relatively low. The concentration required to achieve 50% inhibition (LC50) was 118.16%, indicating that the extract has low toxicity and requires high concentrations for significant inhibition.

REFERENCES

- Anugrah, F. M., & Widiyanti, F. (2018). Pengaruh fungisida berbahan aktif metalaksil, fenamidone, dan dimetomorf terhadap konidia *Peronosclerospora* Spp. Isolat Klaten. *Jurnal Penelitian Saintek*, 23(1), 21-31.
- Rosa, Y. (2021). Aktivitas antijamur ekstrak etanol daun gambir (*Uncaria gambir* Roxb) terhadap *Candida albicans*. *Jurnal Ilmu Kedokteran dan Kesehatan*, 8(3).
- Safitri, K. A., Putra, T. A., & Irawan, A. (2023). Identifikasi Senyawa Metabolit Sekunder Ekstrak Etanolik Kulit Umbi bit (*Beta vulgaris* L.). *Jurnal Ilmiah Bakti Farmasi*, 7(2), 5-9.