

MINFLUX and MINSTED provide molecule-scale resolution in fluorescence microscopy

Stefan W. Hell

Max Planck Institute for Multidisciplinary Sciences, Göttingen &
Max Planck Institute for Medical Research, Heidelberg

I will show how an in-depth description of the basic principles of diffraction-unlimited fluorescence microscopy (nanoscopy) [1] has spawned a new powerful superresolution concept, namely MINFLUX nanoscopy [2-5]. MINFLUX utilizes a local excitation intensity minimum (of a doughnut or a standing wave) that is targeted like a probe in order to localize the fluorescent molecule to be registered. In combination with single-molecule switching, MINFLUX and its more recent ‘cousin’ MINSTED [6] have obtained the ultimate (super)resolution: the size of a molecule. Providing 1–3 nanometer resolution these novel microscopy concepts are being established for routine fluorescence imaging at the highest, molecular-size resolution levels. Relying on fewer detected photons than popular camera-based localization, MINFLUX and MINSTED nanoscopy are poised to open a new chapter in the imaging of protein complexes and distributions in fixed and living cells.

- [1] Hell, S.W. **Nat. Methods** 6, 24-32 (2009).
- [2] Balzarotti, F., Eilers, Y., Gwosch, K. C., Gynnå, A. H., Westphal, V., Stefani, F. D., Elf, J., Hell, S.W. **Science** 355, 606-612 (2017).
- [3] Eilers, Y., Ta, H., Gwosch, K. C., Balzarotti, F., Hell, S. W. **PNAS** 115, 6117-6122 (2018).
- [4] Gwosch, K. C., Pape, J. K., Balzarotti, F., Hoess, P., Ellenberg, J., Ries, J., Hell, S. W. **Nat. Methods** 17, 217-224 (2020)
- [5] Schmidt R., Weihs T., Wurm C., Janssen I., Rehman J., Sahl S.J., Hell S.W. **Nat Commun** 12:1478 (2021)
- [6] Weber, M., Leutenegger M., Stoldt S., Jakobs S., Mihaila T.S., Butkevitch A.N., Hell S.W., **Nat. Photonics**, <https://doi.org/10.1038/s41566-021-00774-2> (2021).