


22.4 MAGNETIC FIELD STRENGTH: FORCE ON A MOVING CHARGE IN A MAGNETIC FIELD

1. What is the direction of the magnetic force on a **positive charge** that moves as shown in each of the six cases shown in Figure 22.56?



2. Repeat Exercise 1 for a **negative charge**.

3. What is the direction of the velocity of a **negative charge** that experiences the magnetic force shown in each of the three cases in Figure 22.57, assuming it moves perpendicular to B ?




Figure 22.57

4. Repeat 3 for a **positive charge**.

5. What is the direction of the magnetic field that produces the magnetic force on a **positive charge** as shown in each of the three cases in the figure below, assuming B is perpendicular to v ?




Figure 22.58

6. Repeat Exercise 5 for a **negative charge**.

7. What is the maximum force on an aluminum rod with a $0.100\text{-}\mu\text{C}$ charge that you pass between the poles of a 1.50-T permanent magnet at a speed of 5.00 m/s ? In what direction is the force?

8. (a) Aircraft sometimes acquire small static charges. Suppose a supersonic jet has a $0.500\text{-}\mu\text{C}$ charge and flies due west at a speed of 660 m/s over the Earth's south magnetic pole, where the $8.00 \times 10^{-5}\text{-T}$ magnetic field points straight up. What are the direction and the magnitude of the magnetic force on the plane? (b) Discuss whether the value obtained in part (a) implies this is a significant or negligible effect.

9. (a) A cosmic ray proton moving toward the Earth at $5.00 \times 10^7\text{ m/s}$ experiences a magnetic force of $1.70 \times 10^{-16}\text{ N}$. What is the strength of the magnetic field if there is a 45° angle between it and the proton's velocity? (b) Is the value obtained in part (a) consistent with the known strength of the Earth's magnetic field on its surface? Discuss.

22.5 FORCE ON A MOVING CHARGE IN A MAGNETIC FIELD: EXAMPLES AND APPLICATIONS

12. A cosmic ray electron moves at 7.50×10^6 m/s perpendicular to the Earth's magnetic field at an altitude where field strength is 1.00×10^{-5} T. What is the radius of the circular path the electron follows?
13. A proton moves at 7.50×10^7 m/s perpendicular to a magnetic field. The field causes the proton to travel in a circular path of radius 0.800 m. What is the field strength?
14. (a) Viewers of Star Trek hear of an antimatter drive on the Starship Enterprise. One possibility for such a futuristic energy source is to store antimatter charged particles in a vacuum chamber, circulating in a magnetic field, and then extract them as needed. Antimatter annihilates with normal matter, producing pure energy. What strength magnetic field is needed to hold antiprotons, moving at 5.00×10^7 m/s in a circular path 2.00 m in radius? Antiprotons have the same mass as protons but the opposite (negative) charge. (b) Is this field strength obtainable with today's technology or is it a futuristic possibility?
15. (a) An oxygen-16 ion with a mass of 2.66×10^{-26} kg travels at 5.00×10^6 m/s perpendicular to a 1.20-T magnetic field, which makes it move in a circular arc with a 0.231-m radius. What positive charge is on the ion? (b) What is the ratio of this charge to the charge of an electron? (c) Discuss why the ratio found in (b) should be an integer.
16. What radius circular path does an electron travel if it moves at the same speed and in the same magnetic field as the proton in Exercise 22.13?

AP workbook

3. A charged particle with constant speed enters a uniform magnetic field whose direction is perpendicular to the particles velocity. The particle will:

- A) Speed up
- B) Experience no change in velocity
- C) Follow a parabolic arc
- D) Follow a circular arc

A long straight wire conductor is placed below a compass as shown in the top view figure.

When a large


conventional current flows in the conductor as shown, the N pole of the compass:

- A) has its polarity reversed
- B) points to the south
- C) points to the west
- D) points to the east




6. Two bar magnets are to be cut in half along the dotted lines shown. None of the pieces are rotated. After the cut:

- A) The two halves of each magnet will attract each other
- B) The two halves of each magnet will repel each other
- C) The two halves of the top magnet will repel, the two halves of the bottom magnet will attract
- D) The two halves of the top magnet will attract, the two halves of the bottom magnet will repel




7. An ion with charge q , mass m , and speed v enters a magnetic field B and is deflected into a path with a radius of curvature R . If a second ion has speed $2v$, while m , q , and B are unchanged, what will be the radius of the second ion's path?

- A) $4R$
- B) $2R$
- C) $R/2$
- D) $R/4$



10. The diagram to the right depicts iron filings sprinkled around three permanent magnets. Pole R is the same pole as

- A) T and Y
- B) T and Z
- C) X and Y
- D) X and Z




14. A positively charged particle moves to the right. It enters a region of space in which there is an electric field directed up the plane of the paper as shown. In which direction does the magnetic field have to point in this region so that the particle maintains a constant velocity?

- A) into the plane of the page
- B) out of the plane of the page
- C) to the right
- D) to the left

23. An electron moves in the plane of the page through two regions of space along the dotted-line trajectory shown in the figure. There is a uniform electric field in Region I directed into the plane of the page (as shown). There is no electric field in Region II. What is a necessary direction of the magnetic field in regions I and II? Ignore gravitational forces.


Region I Region II

- A) Toward bottom of page Up on the page
- B) Toward top of page Into the page
- C) Toward top of page Out of the page
- D) Toward bottom of page Out of the page




29. An electron is in a uniform magnetic field $\mathbf{B}$ that is directed out of the plane of the page, as shown. When the electron is moving in the plane of the page in the direction indicated by the arrow, the force on the electron is directed

- (A) toward the right
- (B) out of the page
- (C) into the page
- (D) toward the top of the page




AP Physics Free Response Practice – Magnetism and Electromagnetism

SECTION A – Magnetism



1975B6. In a mass spectrometer, singly charged ^{16}O ions are first accelerated electrostatically through a voltage V to a speed v_0 . They then enter a region of uniform magnetic field B directed out of the plane of the paper.


- The ^{16}O ions are replaced with singly charged ^{32}S ions of twice the mass and the same charge. What will be their speed in terms of v_0 for the same accelerating voltage?
- When ^{32}S is substituted for ^{16}O in part (a), determine by what factor the radius of curvature of the ions' path in the magnetic field changes.



1976B4.


An ion of mass m and charge of known magnitude q is observed to move in a straight line through a region of space in which a uniform magnetic field B points out of the paper and a uniform electric field E points toward the top edge of the paper, as shown in region I above. The particle travels into region II in which the same magnetic field is present, but the electric field is zero. In region II the ion moves in a circular path as shown.

- Indicate on the diagram below the direction of the force on the ion at point P_2 in region II.



- (b) Is the ion positively or negatively charged? Explain clearly the reasoning on which you base your conclusion.
 (c) Indicate and label clearly on the diagram below the forces which act on the ion at point P_1 in region I.

- (d) Find an expression for the ion's speed v at point P_1 in terms of E and B .



1993B3. A particle of mass m and charge q is accelerated from rest in the plane of the page through a potential difference V between two parallel plates as shown. The particle is injected through a hole in the right-hand plate into a region of space containing a uniform magnetic field of magnitude B oriented perpendicular to the plane of the page. The particle curves in a semicircular path and strikes a detector.

- a. i. State whether the sign of the charge on the particle is positive or negative.
 ii. State whether the direction of the magnetic field is into the page or out of the page.
 b. Determine each of the following in terms of m , q , V , and B .
 i. The speed of the charged particle as it enters the region of the magnetic field B
 ii. The force exerted on the charged particle by the magnetic field B
 iii. The distance from the point of injection to the detector
 iv. The work done by the magnetic field on the charged particle during the semicircular trip

