Engineering an Angular Momentum Bias with Linear Synthetic Motion

João C. Serra, Mário G. Silveirinha* [1]
1. University of Lisbon–Instituto Superior Técnico and Instituto de Telecomunicações, ~ Avenida Rovisco Pais, 1, 1049-001 Lisboa, Portugal

Topological photonics provides an elegant framework to classify different classes of photonic crystals and waveguides with complete band gaps [1]. Conventional photonic crystals are material structures with a periodic spatial modulation of the refractive index. Recently, time has emerged as a new degree of freedom in system design. In particular, the crystal notion has been extended to structures presenting *time* modulations [2-5]. Time and spacetime modulations enable exotic and nonreciprocal light-matter interactions, which can be useful for a myriad of optical applications.

Crystals with a spacetime travelling-wave modulation, $\varepsilon(x - vt, y, z)$ with v the modulation speed are especially significant as they enable a relatively simple analytical modelling and because they can be used to engineer a synthetic translational motion [2, 3]. Heuristically, nontrivial topological phases must be associated with some form of internal angular momentum [6], analogous to the angular momentum imparted on a charged particle by a magnetic field, which is at the origin of the nontrivial topology of gyrotropic materials. This property suggests that travelling-wave spacetime modulations are unsuitable to generate nontrivial topological phases. In this talk, we will show that surprinsingly this understanding is wrong, and that it is possible to engineer a synthetic spacetime rotation and nontrivial topological phases in crystals with a pure travelling-wave (linear) modulation. Our findings are not only interesting from a theoretical standpoint, but also unveil an exciting path to engineer nontrivial material topologies in spacetime crystals.

References

[1] T. Ozawa, H. M. Price, A. Amo, N. Goldman, M. Hafezi, L. Lu, M. C. Rechtsman, D. Schuster, J. Simon, O. Zilberberg, I. Carusotto, "Topological Photonics", *Rev. Mod. Phys.* **91**, 15006 (2019).

[2] Z.-L. Deck-Leger, N. Chamanara, M. Skorobogatiy, M. G. Silveirinha, C. Caloz, "Uniform-velocity spacetime crystals", Adv. Photonics 1, 1 (2019).

[3] P. A. Huidobro, E. Galiffi, S. Guenneau, R. V. Craster, J. B. Pendry, "Fresnel drag in space-time-modulated metamaterials", Proc. Natl. Acad. Sci. 10;116(50):24943-24948 (2019).

[4] E. Galiffi, R. Tirole, S. Yin, H. Li, S. Vezzoli, P. A. Huidobro, M. G. Silveirinha, R. Sapienza, A. Alù, J. B. Pendry, "Photonics of Time-Varying Media", Adv. Photon. 4(1), 014002 (2022), doi: 10.1117/1.AP.4.1.014002.

[5] J. C. Serra, M. G. Silveirinha, "Rotating Spacetime Modulation: Topological Phases and Spacetime Haldane Model", Phys. Rev. B, **107**, 035133, 2023.

[6] M. G. Silveirinha, "Quantized Angular Momentum in Topological Optical Systems", Nat. Comm., **10**, 349, 2019.

Relevant references for talk

J. C. Serra, M. G. Silveirinha, "Rotating Spacetime Modulation: Topological Phases and Spacetime Haldane Model", Phys. Rev. B, 107, 035133, 2023.

F. R. Prudêncio, M. G. Silveirinha, "Ill-defined topologies in local dispersive photonic systems", Phys. Rev. Lett., 129, 133903, 2022.