

G. Álvarez-Herrera, F. Agnolin, et F. Novas. A Rostral Neurovascular System in the Mosasaur *Taniwhasaurus Antarcticus*, 2020.

N. Bardet, A. Houssaye, J-C. Rage, et X. Pereda-Suberbiola. The Cenomanian-Turonian (Late Cretaceous) radiation of marine squamates (Reptilia): The role of the Mediterranean Tethys, 2008.

N. Bardet, A. Houssaye, P. Vincent, X. Pereda-Suberbiola, M. Amaghzaz, E. Jourani et S. Meslouh. Mosasaurids (Squamata) from the Maastrichtian Phosphates of Morocco: Biodiversity, palaeobiogeography and palaeoecology based on tooth morphoguilds, 2015.

P. Brazaitis et M. Watanabe. Crocodilian behaviour: A window to dinosaur behaviour? , 2011.

J. M. Crowe-Riddell, E. P. Snelling, A. P. Watson, A. Kyuseop Suh, J. C. Partridge et K. L. Sanders. The evolution of scale sensilla in the transition from land to sea in elapid snakes, 2016.

M. J. Everhart et K. Ewell. Shark-bitten dinosaur (Hadrosauridae) caudal vertebrae from the Niobrara Chalk (Upper Coniacian) of western Kansas, 2006.

G. Garcia, N. Bardet, A. Houssaye, X. Pereda-Suberbiola et X. Valentin. Mosasauroid (Squamata) discovery in the Late Cretaceous (Early Campanian) continental deposits of Villeveyrac–L’Olivet, southern France, 2015.

D. V. Grigoriev. [Giant Mosasaurus Hoffmanni \(Squamata, Mosasauridae\) from the Late Cretaceous \(Maastrichtian\) of Penza, Russia](#), 2014.

A. Houssaye, J. Lindgren, R. Pellegrini, A. H. Lee, D. Germain et M. J. Polcyn. [Microanatomical and Histological Features in the Long Bones of Mosasaurine Mosasaurs \(Reptilia, Squamata\) – Implications for Aquatic Adaptation and Growth Rates](#), 2013.

L. Kocsis, A. Ősi, T. Vennemann, C. N. Trueman et M. R. Palmer. Geochemical study of vertebrate fossils from the Upper Cretaceous (Santonian) Csehbánya Formation (Hungary): evidence for a freshwater habitat of mosasaurs and pycnodont fish, 2009.

L. Makádi, M. W. Caldwell, et A. Ősi. [The First Freshwater Mosasauroid \(Upper Cretaceous, Hungary\) and a New Clade of Basal Mosasauroids](#), 2012.

D. B. Leitch et K. C. Catania. [Structure, innervation and response properties of integumentary sensory organs in crocodilians](#), 2012.

J. Lindgren, J. W. M. Jaght et M. W. Caldwell. A fishy mosasaur: the axial skeleton of *Plotosaurus* (Reptilia, Squamata) reassessed, 2007.

J. Lindgren, C. Alwmark, M. W. Caldwell et A. R. Fiorillo. [Skin of the Cretaceous mosasaur *Plotosaurus*: implications for aquatic adaptations in giant marine reptiles](#), 2009.

J. Lindgren , H. F. Kaddumi et M. J. Polcyn. Soft tissue preservation in a fossil marine lizard with a bilobed tail fin, 2013.

J. Lindgren, M. J. Polcyn et B. A. Young. Landlubbers to leviathans: evolution swimming in mosasaurine mosasaurs, 2011.

N. R. Longrich, N. Bardet, A. S. Schulp et N-E. Jalil. *Xenodens calminechari* Gen. et Sp. Nov., a Bizarre Mosasaurid (Mosasauridae, Squamata) with Shark-like Cutting Teeth from the Upper Maastrichtian of Morocco, North Africa, 2021a.

N. R. Longrich, N. Bardet, F. Khaldoune, O. Khadiri Yazami et N.-E. Jalil. *Pluridens serpentis*, a new mosasaurid (Mosasauridae: Halisaurinae) from the Maastrichtian of Morocco and implications for mosasaur diversity, 2021b.

T. W. Reeder, T. M. Townsend, D. G. Mulcahy, G. Daniel. B. P. Noonan, P. L. Wood, Perry, J. W. Sites, J. J. Wiens et J. John. [Integrated Analyses Resolve Conflicts over Squamate Reptile Phylogeny and Reveal Unexpected Placements for Fossil Taxa](#), 2015.

A. S. Schulp, E. W. A Mulder et K. Schwenk. Did mosasaurs have forked tongues?, 2005.

T. R. Simões, O. Vernygora, I. Paparella, P. Jimenez-Huidobro et M. W. Caldwell.
[Mosasauroid phylogeny under multiple phylogenetic methods provides new insights on the evolution of aquatic adaptations in the group](#), 2017.