



(510) 679-3300
27206 Calaroga Ave - Ste 201
Hayward, CA 94545
BALMEDS.com

Patient Education

Tadalafil 100mg with Apomorphine 8mg

This guide will help you understand how to safely and effectively use the combination medication **Tadalafil 100mg with Apomorphine 8mg** for managing erectile dysfunction (ED) symptoms.

What is Tadalafil 100mg with Apomorphine 8mg?

This combination therapy includes:

- **Tadalafil (100mg):** A phosphodiesterase type 5 (PDE5) inhibitor that enhances blood flow to the penis, helping to achieve and maintain an erection.
- **Apomorphine (8mg):** A dopamine receptor agonist that stimulates sexual arousal by acting on the central nervous system.

Together, these medications address both the physical and psychological aspects of ED, providing a more comprehensive solution.

How Does it Work?

1. **Tadalafil:** Works by relaxing the blood vessels in the penis, improving blood flow during sexual stimulation.
2. **Apomorphine:** Enhances brain signaling to trigger arousal and improve the erectile response.

This combination works synergistically to help you achieve stronger, longer-lasting erections when sexually stimulated.

How to Take Tadalafil with Apomorphine

1. **Dosage:** Follow your healthcare provider's instructions. The usual dose is one tablet taken as needed, approximately 30-60 minutes before sexual activity.
2. **Administration:** Take the medication on an empty stomach with water to ensure optimal absorption. Avoid high-fat meals, as they may reduce effectiveness.
3. **Do Not Exceed:** Do not take more than one dose in 24 hours.

Possible Side Effects

As with any medication, side effects may occur. Common side effects include:

- **Tadalafil:** Headache, flushing, nasal congestion, upset stomach, back pain, or muscle aches.
- **Apomorphine:** Dizziness, nausea, sweating, or mild drowsiness.

Seek Medical Attention If You Experience any of these rare side effects:

- Severe dizziness or fainting.
- Prolonged or painful erections lasting more than 4 hours (priapism).
- Sudden vision or hearing loss.



(510) 679-3300
27206 Calaroga Ave - Ste 201
Hayward, CA 94545
BALMEDS.com

Precautions and Warnings

1. **Alcohol:** Limit alcohol intake as it may increase the risk of side effects like dizziness and low blood pressure.
2. **Driving or Operating Machinery:** Apomorphine may cause drowsiness or dizziness. Avoid activities requiring alertness if affected.
3. **Pre-existing Conditions:** Inform your doctor if you have any heart conditions, low blood pressure, liver or kidney disease, or a history of priapism.
4. **Drug Interactions:**
 - Do not combine with nitrates or other medications that lower blood pressure.
 - Avoid combining with other PDE5 inhibitors or medications affecting dopamine levels without medical advice.

When to Call Your Doctor

Contact your healthcare provider if you:

- Experience severe side effects or allergic reactions (rash, itching, swelling).
- Have persistent side effects interfering with daily life.
- Notice that the medication is not working as expected.

Storage

- Store at room temperature (20-25°C or 68-77°F).
- Keep away from moisture, heat, and direct sunlight.
- Keep out of reach of children and pets.

Additional Tips for Success

- Engage in foreplay to maximize the medication's effects.
- Maintain a healthy lifestyle, including regular exercise, balanced nutrition, and stress management, to improve overall erectile function.

If you have any further questions, consult your healthcare provider for personalized advice.

It is important that you continue to see your regular medical doctor for your usual health care, tests, and routine health maintenance. We are not replacing your primary medical doctor.

● ————— ● **#HelloHappiness** ● ————— ●