

Gastric carcinoma during pregnancy: Report of a case

Gebelikte saptanan mide kanseri: Olgu sunumu

İlkay Tuğba ÜNEK, Aygül ÇELTİK, Ahmet ALACACIOĞLU, Suna ÇOKMERT, Tuğba YAVUZŞEN, Nedim Serkan DOĞAN, İlhan ÖZTOP, Binnaz DEMIRKAN, Uğur YILMAZ

Department of Internal Medicine, Division of Medical Oncology, Dokuz Eylül University, School of Medicine, Izmir

Background/aims: Gastric cancer in pregnancy is extremely rare and often diagnosed at advanced stages. Well-recognized pregnancy-related symptoms, such as nausea and epigastric discomfort, can be the first symptoms of gastric cancer. Thus, the diagnosis of gastric cancer in pregnancy is difficult. We herein report a case of gastric cancer in pregnancy to alert clinicians to this rare possibility. **Materials and methods:** A 22-year-old nulliparous woman was examined by a gynecologist with a complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. Gastroscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells and her pregnancy was terminated. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. Tumor invasion was detected in pancreas and spleen. She was admitted to the medical oncology clinic and received a palliative chemotherapy. Bilateral double-J-stent placement was performed because of obstructive uropathy. A nasogastric tube was inserted for intestinal decompression, and total parenteral nutrition was administered. An adequate pain medication was given. To date (3 months after the diagnosis) the patient has been well without any signs of progression. **Conclusion:** Early diagnosis of gastric cancer is very important for a better outcome. The diagnosis may be delayed because mild gastrointestinal symptoms are common during pregnancy. Clinicians should take this into consideration in the differential diagnosis of persistent epigastric complaint during pregnancy. Suspicion and early upper gastrointestinal endoscopy are necessary.

Key words: Gastric cancer, pregnancy, chemotherapy

INTRODUCTION

Gastric cancer during pregnancy is extremely rare, with a reported incidence of 0.1% (1). This reflects its tendency to manifest with advancing age and its male preponderance (2). The diagnosis of gastric cancer in pregnancy is very often delayed.

Amacı: Gebelikte mide kanseri çok nadirdir ve sıklıkla ileri evrelerde tanı koyulur. Bulantı, epigastrik rahatsızlık gibi gebelikte ilişkili olduğu iyi bilinen semptomlar, mide kanserinin ilk semptomları olabilir. Bu nedenle, gebelikte mide kanseri tanısı zordur. Burada, bu nadir olasığa klinisyenlerin dikkatini çekmek amacıyla, gebelikte mide kanseri tanısı almış bir olgu sunulmaktadır. **Yöntem:** 22 yaşında kadın hasta kasık ağrısı şikayetiyle bir jinekolog tarafından muayene edildi. Karın ultrasonografisinde 7 haftalık gebe olduğu saptandı ve ayrıca yoğun miktarda asit görüldü. Hasta daha ileri değerlendirme amacıyla bir hastaneye yönlendirildi. Hastanın iştahsızlığı dışında iştahsızlık, erken doyma ve yemek sonrası şişkinlik şikayetleri vardı. Yapılan gastroskopiye, kardiyadan küken alan ve tüm distal korpusu invaze eden tümör saptandı. Histopatolojik incelemede taşlı yüzük hücreli mide karsinomu tanısı koyuldu ve hastanın gebeliği sonlandırıldı. Exploratif laparotomide unresektabl mide kanseri ve multiple peritoneal implantlar saptandı. Tümörün, pankreas ve dalaga invaze olduğu görüldü. Tıbbi onkoloji kliniğine kabul edilen hastaya palyatif kemoterapi verildi. Obstrüktif üropati nedeniyle bilateral double-J-stent yerleştirildi. İntestinal dekompresyon için nazogastrik sonda takıldı ve total parenteral beslenme uygulandı. Yeterli ağrı palyasyonu yapıldı. Şimdiye kadar (tanıdan 3 ay sonrasında) herhangi bir progresyon bulgusu gözlenmeyen hastanın durumu iyidir. **Sonuç:** Mide kanserinin erken tanısı, daha iyi sonuçlar elde edilmesi açısından çok önemlidir. Tanıda gecikmeler olabilir çünkü hafif gastrointestinal semptomlar gebelikte sık görülür. Klinisyenler, gebelikte sürekli olan epigastrik şikayetlerin ayırıcı tanısında mide kanserini de düşünmelidir. Tanı için, mide kanserinden şüphe duyulması ve erken gastroskopi yapılması gereklidir.

Anahtar kelimeler: Mide kanseri, gebelik, kemoterapi

In 97% of the cases, patients were found to be at advanced tumor stage when curative therapies are not possible (3). Its prognosis is generally poor. Early diagnosis of gastric cancer is very important for a better outcome (3). The diagnosis of

Address for correspondence: İlkay Tuğba ÜNEK
Department of Internal Medicine, Division of Medical Oncology
Dokuz Eylül University, School of Medicine, 35340, Inciraltı, Izmir
Phone: +90 232 412 48 01 • Fax: +90 232 412 37 55
E-mail: tugba.unek@deu.edu.tr

Manuscript received: 27.07.2006 Accepted: 01.02.2007

Gastric carcinoma during pregnancy: Report of a case

Gebelikte saptanan mide kanseri: Olgu sunumu

İlkay Tuğba ÜNEK, Aygül ÇELTİK, Ahmet ALACACIOĞLU, Suna ÇOKMERT, Tuğba YAVUZFİEN, Nedim Serkan DOĞAN, İlhan ÖZTOP, Binnaz DEMİRKAN, Uğur YILMAZ

Department of Internal Medicine, Division of Medical Oncology, Dokuz Eylül University, School of Medicine, İzmir

Background/aims: Gastric cancer in pregnancy is extremely

rare and often diagnosed at advanced stages. Well-recognized

gestation-related symptoms, such as nausea and epigastric

discomfort, can be the first symptoms of gastric cancer. Thus,

the diagnosis of gastric cancer in pregnancy is difficult. We here-

in report a case of gastric cancer in pregnancy to alert clinicians

to this rare possibility. Materials and methods: A 22-year-old nulliparous woman was examined by a gynecologist with a complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. Gastroscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells and her pregnancy was terminated. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. Tumoral invasion was detected in pancreas and spleen. She was admitted to the medical oncology clinic and received a palliative chemotherapy. Bilateral double-J-stent placement was performed because of obstructive uropathy. A nasogastric tube was inserted for intestinal decompression, and total parenteral nutrition was administered. An adequate pain medication was given. To date (3 months after the diagnosis) the patient has been well without any signs of progression. Conclusion: Early diagnosis of gastric

Address for correspondence: İlkay Tuğba ÜNEK Manuscript received: 27.07.2006 Accepted: 01.02.2007 Department of Internal Medicine, Division of Medical Oncology Dokuz Eylül University, School of Medicine, 35340, İzmir Phone: +90 232 412 48 01 • Fax: +90 232 412 37 55 E-mail: tugba.gun@deu.edu.tr

sunulmaktadır. Yöntem: 22 yaşında kadın hasta kasıksız şikayetiyle bir jinekolog tarafından muayene edildi. Karın ultrasonografisinde 7 haftalık gebelik saptandı ve ayrıca yoğun miktarda asit görüldü. Hasta daha ileri değerlendirme amacıyla bir hastaneye yönlendirildi. Hastanın kasıksız şikayeti, erken doyma ve yemek sonrası şişkinlik şikayetleri vardı. Yapılan gastroskopi, kardiyadan köken alan ve tüm distal korpusu invaze eden tümör saptandı. Histopatolojik incelemede taflı yüzük hücreli mide karsinomu tanısı koyuldu ve hastanın gebeliği sonlandırıldı. Exploratif laparotomide rezektabl mide kanseri ve multiple peritoneal implantlar saptandı. Tümörün, pankreasa ve dalaca invaze olduğu görüldü. Tıbbi onkoloji kliniğine kabul edilen hastaya palyatif kemoterapi verildi. Obstrüktif üropati nedeniyle bilateral double-J-stent yerleştirildi. (intestinal dekompresyon için nazogastrik sonda takıldı) ve total parenteral beslenme uygulandı. Yeterli palyasyonu yapıldı. şimdiye kadar (tanıdan 3 ay sonrasında) herhangi bir progresyon bulgusu gözlenmeyen hastanın durumu iyidir. Sonuç: Mide kanserinin erken tanısı, ric cancer is very important for a better outcome. The diagnosis

may be delayed because mild gastrointestinal symptoms are

Amaç: Gebelikte mide kanseri çok nadirdir ve sıklıkla ileri

evrelerde tanı koyulur. Bulantı, epigastrik rahatsızlık gibi

ile ilişkili olduğu iyi bilinen semptomlar, mide kanserinin

semptomları olabilir. Bu nedenle, gebelikte mide kanseri

zordur. Burada, bu nadir olası bir klinisyenlerin dikkatini

emek amacıyla, gebelikte mide kanseri tanısı almış bir olgu

ar-old nulliparous woman was examined by a gynecologist with a

complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed

massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. Gastroscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells and her pregnancy was terminated. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. Tumoral invasion was detected in pancreas and spleen. She was admitted to the medical oncology clinic and received a palliative chemotherapy. Bilateral double-J-stent placement was performed because of obstructive uropathy. A nasogastric tube was inserted for intestinal decompression, and total parenteral nutrition was administered. An adequate pain medication was given. To date (3 months after the diagnosis) the patient has been well without any signs of progression. Conclusion: Early diagnosis of gastric

Address for correspondence: İlkay Tuğba ÜNEK Manuscript received: 27.07.2006 Accepted: 01.02.2007 Department of Internal Medicine, Division of Medical Oncology Dokuz Eylül University, School of Medicine, 35340, İzmir Phone: +90 232 412 48 01 • Fax: +90 232 412 37 55 E-mail: tugba.gun@deu.edu.tr

sunulmaktadır. Yöntem: 22 yaşında kadın hasta kasıksız şikayetiyle bir jinekolog tarafından muayene edildi. Karın ultrasonografisinde 7 haftalık gebelik saptandı ve ayrıca yoğun miktarda asit görüldü. Hasta daha ileri değerlendirme amacıyla bir hastaneye yönlendirildi. Hastanın kasıksız şikayeti, erken doyma ve yemek sonrası şişkinlik şikayetleri vardı. Yapılan gastroskopi, kardiyadan köken alan ve tüm distal korpusu invaze eden tümör saptandı. Histopatolojik incelemede taflı yüzük hücreli mide karsinomu tanısı koyuldu ve hastanın gebeliği sonlandırıldı. Exploratif laparotomide rezektabl mide kanseri ve multiple peritoneal implantlar saptandı. Tümörün, pankreasa ve dalaca invaze olduğu görüldü. Tıbbi onkoloji kliniğine kabul edilen hastaya palyatif kemoterapi verildi. Obstrüktif üropati nedeniyle bilateral double-J-stent yerleştirildi. (intestinal dekompresyon için nazogastrik sonda takıldı) ve total parenteral beslenme uygulandı. Yeterli palyasyonu yapıldı. şimdiye kadar (tanıdan 3 ay sonrasında) herhangi bir progresyon bulgusu gözlenmeyen hastanın durumu iyidir. Sonuç: Mide kanserinin erken tanısı, ric cancer is very important for a better outcome. The diagnosis

may be delayed because mild gastrointestinal symptoms are

Amaç: Gebelikte mide kanseri çok nadirdir ve sıklıkla ileri

evrelerde tanı koyulur. Bulantı, epigastrik rahatsızlık gibi

ile ilişkili olduğu iyi bilinen semptomlar, mide kanserinin

semptomları olabilir. Bu nedenle, gebelikte mide kanseri

zordur. Burada, bu nadir olası bir klinisyenlerin dikkatini

emek amacıyla, gebelikte mide kanseri tanısı almış bir olgu

ar-old nulliparous woman was examined by a gynecologist with a

complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. Gastroscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells and her pregnancy was terminated. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. Tumoral invasion was detected in pancreas and spleen. She was admitted to the medical oncology clinic and received a palliative chemotherapy. Bilateral double-J-stent placement was performed because of obstructive uropathy. A nasogastric tube was inserted for intestinal decompression, and total parenteral nutrition was administered. An adequate pain medication was given. To date (3 months after the diagnosis) the patient has been well without any signs of progression. Conclusion: Early diagnosis of gastric

semp- common during pregnancy. Clinicians should take this into

olan consideration in the differential diagnosis of persistent epigast-

ric complaint during pregnancy. Suspicion and early upper

ve gastrointestinal endoscopy are necessary.

erken gastroskopi yapılmas› gereklidir.

Key words: Gastric cancer, pregnancy, chemotherapy

Anahtar kelimeler: Mide kanseri, gebelik, kemoterapi

INTRODUCTION

Gastric cancer during pregnancy is extremely rare, with a reported incidence of 0.1% (1). This reflects its tendency to manifest with advancing age and its male preponderance (2). The diagnosis of gastric cancer in pregnancy is very often delayed.

In 97% of the cases, patients were found to be at advanced tumor stage when curative therapies are not possible (3). Its prognosis is generally poor. Early diagnosis of gastric cancer is very important for a better outcome (3). The diagnosis of

Tan›da gecikmeler olabilir çünkü hafif gastrointestinal

tomlar gebelikte sık görülür. Klinisyenler, gebelikte sürekli

epigastrik şikayetlerin ay›r›c› tan›s›nda mide kanserini de

düflünmelidir. Tan› için, mide kanserinden flüphe duyulmas›

gastric cancer in pregnancy is difficult, because early symptoms will usually be misinterpreted as the common pregnancy-induced nausea, mild epigastric pain, fullness, etc. (2). We herein report a patient who suffered from advanced gastric cancer during the first trimester of pregnancy, to alert clinicians to this rare possibility. The differential diagnosis of persistent epigastric discomfort associated with pregnancy must include gastric cancer. Suspicion and early upper gastrointestinal endoscopy are necessary for the diagnosis (2).

CASE REPORT

A 22-year-old nulliparous woman was examined by a gynecologist with a complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. There was a history of gastric cancer in her grandfather. Upper gastrointestinal endoscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells, and her pregnancy was terminated. Computerized tomography of the abdomen showed a huge mass in the stomach, massive ascites and retroperitoneal lymphadenopathies. Tumoral invasion was detected in pancreas and spleen. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. She was subsequently admitted to the medical oncology clinic for systemic chemotherapy.

On admission, the patient was experiencing significant abdominal pain, general malaise, nausea, vomiting, and constipation, and was unable to tolerate oral intake. On physical examination, we found a cachectic patient with a prominent abdomen, epigastric tenderness and ascites. In laboratory examinations, a normocytic anemia with thrombocytosis (hemoglobin 9 g/dl, platelet 600,000/ μ L, and white blood cell count 7,600/ μ L) was detected. Renal function tests were compatible with acute renal failure: blood urea nitrogen 114 mg/dl, serum creatinine 9.31 mg/dl, serum sodium 129 mmol/L, and potassium 5.33 mmol/L. Liver function tests were normal except for hypoalbuminemia (serum albumin 2.5 g/dl). Abdominopelvic ultrasonography showed bilateral dilation of the urinary tract. A double-J-stent placement was per-

formed because of the obstructive uropathy. Her creatinine levels decreased to normal in seven days. The upright abdominal radiograph of this patient revealed stepladder air-fluid levels distributed throughout the abdomen. Based on the diagnosis of an acute mechanical small bowel obstruction, a nasogastric tube was inserted for intestinal decompression, fluid and electrolyte replacement was performed, and total parenteral nutrition was administered through a central venous catheter.

The patient received supportive treatment, adequate pain medication and a palliative chemotherapy. The chemotherapy regimen comprised a 2-h infusion of folinic acid (400 mg m^{-2}), then a bolus injection of 5-fluorouracil (400 mg m^{-2}) on day 1, followed by a 46h infusion of 5-fluorouracil (2400 mg m^{-2}). Epirubicin (75 mg m^{-2}) was administered on day 1 as a 30-min infusion. Cisplatin was not administered because of renal impairment. The regimen was repeated every two weeks. To date (3 months after the diagnosis), the patient has been well without any signs of progression.

DISCUSSION

A case of a 22-year-old woman at the 7th week of gestation who suffered from advanced gastric carcinoma is herein presented. Gastric cancer during pregnancy is extremely rare. It was first reported by Fujimura and Fukuda in 1916 (4). In 1998, Fazeny and Marosi (5) conducted a 30-year review of the world's medical literature. Among 100 reports of pregnancy-associated gastric cancer, only one survivor was detected. Maternal survival of 9-19 months after diagnosis in pregnancy was observed for all other cases. The predominance of a far advanced disease characterized by poorly differentiated adenocarcinomas with presence of peritoneal metastases, as in our case, was detected.

While in older patients the majority of carcinomas are of the intestinal, usually well differentiated type, the tumors in young patients are mainly poorly differentiated carcinomas of the diffuse type with signet ring cells and peritoneal metastasis. These characteristics were more pronounced in the pregnancy-associated cases (6). In our patient, the histopathological diagnosis was also gastric carcinoma with signet ring cells.

The pathogenesis of relationship of pregnancy and gastric cancer remains idiopathic. Besides environmental and genetic etiology, factors associated with the pregnancy itself are considered in the

gastric cancer in pregnancy is difficult, because early symptoms will usually be misinterpreted as the common pregnancy-induced nausea, mild epigastric pain, fullness, etc. (2). We herein report a patient who suffered from advanced gastric cancer during the first trimester of pregnancy, to alert clinicians to this rare possibility. The differential diagnosis of persistent epigastric discomfort associated with pregnancy must include gastric cancer. Suspicion and early upper gastrointestinal endoscopy are necessary for the diagnosis (2).

CASE REPORT

A 22-year-old nulliparous woman was examined by a gynecologist with a complaint of low abdominal pain. An abdominal ultrasound of the patient revealed 7th week of gestation and also showed massive ascites. She was referred to a hospital for further evaluation. Her other symptoms included loss of appetite, early satiety, and postprandial fullness. There was a history of gastric cancer in her grandfather. Upper gastrointestinal endoscopy demonstrated a tumor originating from the cardia and invading throughout the distal corpus. The histopathological diagnosis was gastric carcinoma with signet ring cells, and her pregnancy was terminated. Computerized tomography of the abdomen showed a huge mass in the stomach, massive ascites and retroperitoneal lymphadenopathies. Tumoral invasion was detected in pancreas and spleen. An explorative laparotomy revealed an unresectable gastric cancer and multiple peritoneal implants. She was subsequently admitted to the medical oncology clinic for systemic chemotherapy.

On admission, the patient was experiencing significant abdominal pain, general malaise, nausea, vomiting, and constipation, and was unable to tolerate oral intake. On physical examination, we found a cachectic patient with a prominent abdomen, epigastric tenderness and ascites. In laboratory examinations, a normocytic anemia with thrombocytosis (hemoglobin 9 g/dl, platelet 600,000/ μ L, and white blood cell count 7,600/ μ L) was detected. Renal function tests were compatible with acute renal failure: blood urea nitrogen 114 mg/dl, serum creatinine 9.31 mg/dl, serum sodium 129 mmol/L, and potassium 5.33 mmol/L. Liver function tests were normal except for hypoalbuminemia (serum albumin 2.5 g/dl). Abdominopelvic ultrasonography showed bilateral dilation of the urinary tract. A double-J-stent placement was per-

formed because of the obstructive uropathy. Her creatinine levels decreased to normal in seven days. The upright abdominal radiograph of this patient revealed stepladder air-fluid levels distributed throughout the abdomen. Based on the diagnosis of an acute mechanical small bowel obstruction, a nasogastric tube was inserted for intestinal decompression, fluid and electrolyte replacement was performed, and total parenteral nutrition was administered through a central venous catheter.

The patient received supportive treatment, adequate pain medication and a palliative chemotherapy. The chemotherapy regimen comprised a 2-h infusion of folinic acid (400 mg m^{-2}), then a bolus injection of 5-fluorouracil (400 mg m^{-2}) on day 1, followed by a 46h infusion of 5-fluorouracil (2400 mg m^{-2}). Epirubicin (75 mg m^{-2}) was administered on day 1 as a 30-min infusion. Cisplatin was not administered because of renal impairment. The regimen was repeated every two weeks. To date (3 months after the diagnosis), the patient has been well without any signs of progression.

DISCUSSION

A case of a 22-year-old woman at the 7th week of gestation who suffered from advanced gastric carcinoma is herein presented. Gastric cancer during pregnancy is extremely rare. It was first reported by Fujimura and Fukunda in 1916 (4). In 1998, Fazeny and Marosi (5) conducted a 30-year review of the world's medical literature. Among 100 reports of pregnancy-associated gastric cancer, only one survivor was detected. Maternal survival of 9-19 months after diagnosis in pregnancy was observed for all other cases. The predominance of a far advanced disease characterized by poorly differentiated adenocarcinomas with presence of peritoneal metastases, as in our case,

was detected.

While in older patients the majority of carcinomas are of the intestinal, usually well differentiated type, the tumors in young patients are mainly poorly differentiated carcinomas of the diffuse type with signet ring cells and peritoneal metastasis. These characteristics were more pronounced in the pregnancy-associated cases (6). In our patient, the histopathological diagnosis was also gastric carcinoma with signet ring cells.

The pathogenesis of relationship of pregnancy and gastric cancer remains idiopathic. Besides environmental and genetic etiology, factors associated with the pregnancy itself are considered in the

pathogenesis. Furukawa et al. (7) reported that pregnancy in young females might accelerate the growth of gastric cancer. The promoter effect of sex hormones on carcinogenesis was demonstrated experimentally in the stomach of rats. It was reported that the development and growth of gastric cancer were enhanced by changes in the hormonal circumstances associated with pregnancy (5). However, the effect of estrogen and the results of anti-estrogen treatment in clinical trials are controversial. Jaspers et al. (2) reported that the features and prognosis in gastric cancer associated with pregnancy were the same as in other young patients. Currently, whether or not pregnancy accelerates gastric neoplasia is unknown.

Endoscopic examination has been reported to be safe in pregnancy. Risk factors which were present in most of the reported cases might be helpful for the indication of upper gastrointestinal endoscopy. These include a family history of gastric cancer, a personal history of gastric or duodenal ulcer, immunosuppressive diseases or drugs, and severe smoking (2). In our patient, there was a history of gastric cancer in her grandfather and she had also been suffering from early satiety, postprandial fullness and loss of appetite for several months, which necessitated the urgent upper gastrointestinal endoscopy for early diagnosis of gastric cancer.

Chronic infection with *Helicobacter pylori* (*Hp*) is associated with an elevated risk of developing gastric carcinoma (8). For the diagnosis of the infection in primary care, it is strongly recommended that urea breath test or stool antigen test be

used. Screening for *Hp* inside and outside of pregnancy is only advisable for dyspeptic patients. Patients with alarm symptoms and signs require prompt endoscopic investigation (9). Unfortunately, the *Hp* status of our patient could not be assessed because upper gastrointestinal endoscopy was not performed in our hospital and the patient refused a second endoscopic investigation.

In any case of gastric cancer in pregnancy, a patient's management should be planned according to the guidelines summarized by Ueo et al. (3). In early pregnancy, termination must be discussed to allow optimal treatment. As in this case, in advanced gastric cancer diagnosed in the first trimester, the standard approach is palliative chemotherapy following termination of pregnancy. Thus, herein, the patient's pregnancy was terminated before the chemotherapy. Beyond 24 weeks of gestation, the decision depends on the stage of gastric cancer. If the cancer is advanced and considered resectable, immediate resection of the gastric cancer is recommended despite the risk to the fetus. After week 30, the fetus should be delivered followed by radical operation on the gastric cancer (3).

In conclusion, the only means of possibly improving the prognosis of gastric cancer in pregnancy is detect it at earlier stages. Although rare under the age of 30, clinicians should take this into consideration in the differential diagnosis of persistent epigastric complaint during pregnancy. An extensive diagnostic workup including upper gastrointestinal endoscopy is advisable in patients at risk before empiric treatment with antacid medications.

REFERENCES

1. Silverberg E, Lubera J. Cancer statistics, 1989. *Cancer* 1989; 39: 3-20.
2. Jaspers VKI, Gillesen A, Quakernack K. Gastric cancer in pregnancy: Do pregnancy, age or female sex alter the prognosis? Case reports and review. *Eur J Obstet Gynecol Reprod Biol* 1999; 87: 13-22.
3. Ueo H, Matsuoka H, Tamura S, et al. Prognosis in gastric cancer associated with pregnancy. *World J Surg* 1991; 15: 293-8.
4. Fujimura M, Fukuda K. Gastric cancer associated with pregnancy. *J Kinki Obstet* 1916; 3: 208.
5. Fazeny B, Marosi C. Gastric cancer as an essential differential diagnosis of minor epigastric discomfort during pregnancy. *Acta Obstet Gynecol Scand* 1998; 77: 469-71.
6. Meata M, Yamashiro H, Oka A, et al. Gastric cancer in the young, with special reference to 14 pregnancy-associated cases: analysis based on 2325 consecutive cases of gastric cancer. *J Surg Oncol* 1995; 58: 191-5.
7. Furukawa H, Iwanaga T, Hiratsuka M, et al. Gastric cancer in young adults: growth accelerating effect of pregnancy and delivery. *J Surg Oncol* 1994; 55: 3-6.
8. Meining A, Bayerdorffler E, Muller P, et al. Gastric carcinoma risk index in patients infected with *Helicobacter pylori*. *Virchows Arch* 1998; 432: 311-4.
9. Mulfertheiner P, Megraud F, O'Morain C, et al. Current concepts in the management of *Helicobacter pylori* infection - The Manstricht 2-2000 Consensus Report. *Aliment Pharmacol Ther* 2002; 16: 167-80.

pathogenesis. Furukawa et al. (7) reported that pregnancy in young females might accelerate the growth of gastric cancer. The promoter effect of hormones on carcinogenesis was demonstrated experimentally in the stomach of rats. It was reported that the development and growth of gastric cancer were enhanced by changes in the hormonal circumstances associated with pregnancy

However, the effect of estrogen and the results of anti-estrogen treatment in clinical trials are controversial. Jaspers et al. (2) reported that the features and prognosis in gastric cancer associated with pregnancy were the same as in other young patients. Currently, whether or not pregnancy accelerates gastric neoplasia is unknown.

REFERENCES 1. Silverberg E, Lubera J. Cancer statistics, 1989. *Cancer*

1989; 39: 3-20. 2. Jaspers VKI, Gillessen A, Quakernack K. Gastric cancer in pregnancy: Do pregnancy, age or female sex alter the prognosis? Case reports and review. *Eur J Obstet Gynecol Reprod Biol* 1999; 87: 13-22. 3. Ueo H, Matsuoka H, Tamura S, et al. Prognosis in gastric cancer associated with pregnancy. *World J Surg* 1991; 15: 293-8. 4. Fujimura M, Fukunda K. Gastric cancer associated with

pregnancy. *J Kinki Obstet* 1916; 3: 208. 5. Fazeny B, Marosi C. Gastric cancer as an essential differential diagnosis of minor epigastric discomfort during pregnancy. *Acta Obstet Gynecol Scand* 1998; 77: 469-71.

In any case of gastric cancer in pregnancy, a patient's management should be planned according to the guidelines summarized by Ueo et al. (3). In early pregnancy, termination must be discussed to allow optimal treatment. As in this case, in advanced gastric cancer diagnosed in the first trimester, the standard approach is palliative chemotherapy. Endoscopic examination has been reported to be following termination of pregnancy. Thus, herein, safe the patient's pregnancy was terminated before the sent chemotherapy. Beyond 24 weeks of gestation, the for decision depends on the stage of gastric cancer. If the cancer is advanced and considered resectable, ric

used. Screening for Hp inside and outside of pregnancy is only advisable for dyspeptic patients. Patients with alarm symptoms and signs require sex prompt endoscopic investigation (9). Unfortunately, the Hp status of our patient could not be assessed because upper gastrointestinal endoscopic was not performed in our hospital and the patient

refused a second endoscopic investigation. (5).

cancer, a personal history of gastric or duodenal ulcer, immunosuppressive diseases or drugs, severe smoking (2). In our patient, there was history of gastric cancer in her grandfather and

been suffering from early satiety, postprandial fullness and loss of appetite for several months, which necessitated the urgent upper gastrointestinal endoscopy for early diagnosis of gastric cancer.

In conclusion, the only means of possibly improving the prognosis of gastric cancer in pregnancy is detect it at earlier stages. Although rare under the age of 30, clinicians should take this into consideration in the differential diagnosis of persistent epigastric complaint during pregnancy. An extensive diagnostic workup including upper gastrointestinal endoscopy is advisable in patients at risk before empiric treatment with antacid medications. Chronic infection with *Helicobacter pylori* (Hp) is associated with an elevated risk of developing gastric carcinoma (8). For the diagnosis of the infection in primary care, it is strongly recommended that urea breath test or stool antigen test be

immediate resection of the gastric cancer is recommended despite the risk to the fetus. After week and 30, the fetus should be delivered followed by radi-

cal operation on the gastric cancer (3). she had also

cal operation on the gastric cancer (3). she had also

6. Meata M, Yamashiro H, Oka A, et al. Gastric cancer in the young, with special reference to 14 pregnancy-associated cases: analysis based on 2325 consecutive cases of gastric cancer. *J Surg Oncol* 1995; 58: 191-5. 7. Furukawa H, Iwanaga T, Hiratsuka M, et al. Gastric cancer in young adults: growth accelerating effect of pregnancy and delivery. *J Surg Oncol* 1994; 55: 3-6. 8. Meining A, Bayerdorffer E, Muller P, et al. Gastric carcinoma risk index in patients infected with *Helicobacter pylori*. *Virchows Arch* 1998; 432: 311-4. 9. Malfertheiner P, Megraud F, O'Morain C, et al. Current concepts in the management of *Helicobacter pylori* infection - The Maastricht 2-2000 Consensus Report. *Aliment Pharmacol Ther* 2002; 16: 167-80.