



A title should be the fewest possible words that accurately describe the content of the paper, no more than 16 words (Times New Roman, center, bold, 16pt)

Author 1*, Author 2, Author 3, Author 4, Author 5 (12pt, bold, maximum consisting of 5 authors) 1 Name of University, Country (12pt, Times New Roman)

Correspondence: author@author.com

Article History:

Received date : YYMMDD
Revised date : YYMMDD
Accepted date : YYMMDD
Published date : YYMMDD



Copyright: © 2024 bv the authors. Submitted for possible open access publication under the terms and conditions the Creative Commons Attribution (BY NC) license (https://creativecomm ons.org/licenses/by-n c/4.0/).

ABSTRACT

A well-prepared abstract allows readers to quickly and accurately identify the basic content of a document, determine its relevance to their interests, and thereby decide whether to read the document in its entirety. The abstract must be informative and clear enough, written clearly, and provide a clear statement of the problem, research objectives, research findings, and conclusions. Abstracts should consist of 100 to 200 words. The abstract must be written in the past tense. Standard nomenclature should be used, and abbreviations should be avoided. No literature may be cited. Keyword lists provide the opportunity to add keywords used by indexing and abstracting services in addition to the keywords already present in the title. Wise use of keywords can increase the ease with

which interested parties find our articles (12pt).

Keyword: The first keyword; the second keyword; the third keyword; The fourth keyword; The fifth keywords. (There are a minimum of five keywords and a maximum of six keywords)

INTRODUCTION (Capital, bold, Times new romance 12 pt)

This section explains the three main components. First, to describe the phenomenon being studied, the introduction must contain the research background and research context. Second, the author explains the relationship between the phenomenon and existing theories (at least the journal cited must be less than ten years old), along with gap analysis and the novelty of the research, and finally explains the research objectives. All introductions should be presented in paragraph form, not pointers, with a proportion of 15-20% of the overall length of the article.

The introduction should not be divided into background sub-chapters, problem formulation, and objectives. Beginning of paragraph once tab. Citations are written in bodynote format and are relevant to the bibliography (recommended using the Mendeley application or other reference management application programs such as EndNote, Reference Manager, or Zotero) (12pt, spacing 1.5, spacing after paragraph 6pt).

The manuscript should be written as concisely, consistently, and as directly as possible. The number of pages consists of 10–20 (twenty) pages (including figures and tables). Manuscripts are written single-spaced on one side of A4-sized paper (210 x 297 mm). Manuscripts must have normal margins, or top, bottom, right, and left margins, namely 2.54 cm. The font used is Times New Roman. 12pt. Manuscripts must be written in English.

METHODS

The Methods section must be short but must include sufficient technical information and contain the type of research, research population, research samples or subjects, and data analysis techniques. Only new methods have to be described in detail. Cite previously published procedures in References.

Table 1. Search Strategy

| Database | Search Strategy | Hits |
|-------------------|---------------------------|------|
| Pubmed | ("acne" AND "microbiome") | 7 |
| Science Direct | ("acne" AND "microbiome") | 79 |
| Sagepub | ("acne" AND "microbiome") | 17 |




Figure 1. Article search flow chart

RESULT

Results should include the rationale or design of the experiment as well as the results of the experiment. Results can be presented in the form of images, tables, and text. Research findings must be supported by adequate data. This section must answer the research hypothesis.

Avoid writing in the form of bullet numbering or item list style; it is best to write it in the form of a descriptive paragraph, even though it is a list item. If it contains tables and figures, the numbering is a continuation of the previous number. Each table and figure must be given a title.

Table

The table is in the middle. Use Times New Roman and font sizes 8 to 11. Horizontal lines in the middle of the table do not need to be displayed; only display the heading and the very end, and there should also be no vertical lines. Make sure you create the table correctly via the Insert Table menu. Tables should be referenced in the text by writing something like: '... (Tables are written with a capital 'T').

Table 3. The literature included in this study

| Schneider, et al. ¹¹ (2023) | USA | Cross sectional study | 48 patients | A significant shift in microbial diversity emerged between early (T1-T2) and late (T3-T5) stages of puberty, coinciding with increased sebum production on the face. The overall relative abundance of C. acnes in both normal and acne skin increased during puberty and individual C. acnes strains were uniquely affected by pubertal stage and the presence of acne. Further, an acne microbiome signature associated with unique C. acnes strain composition and metabolic activity emerges in late puberty in those with acne. This unique C. acnes strain composition is predicted to have increased porphyrin production, which may contribute to skin inflammation. |
|--|-------|-----------------------------|--------------|--|
| Cavallo et al. ¹² (2022) | Italy | Cross sectional study | 100 patients | Microbiota analysis showed a significantly lower alpha diversity in inflammatory lesions (LA) than in non-inflammatory |

| | | | | 277.1 |
|----------------------------|-----------|-----------------------------|-------------|-----------------------------|
| | | | | (NI) lesions of acne |
| | | | | patients and healthy |
| | | | | subjects (HS). Differences |
| | | | | at the species level were |
| | | | | driven by the |
| | | | | overabundance of C. acnes |
| | | | | on LA than NI and HS. |
| | | | | The phylotype IA1 was |
| | | | | more represented in the |
| | | | | skin of acne patients than |
| | | | | in HS. Genes involved in |
| | | | | lipids transport and |
| | | | | metabolism, as well as |
| | | | | potential virulence factors |
| | | | | associated with host-tissue |
| | | | | colonization, were detected |
| | | | | in all IA1 strains |
| | | | | independently from the site |
| | | | | of isolation. Additionally, |
| | | | | the IA1 isolates were more |
| | | | | efficient in early adhesion |
| | | | | and biomass production |
| | | | | than other phylotypes |
| | | | | showing a significant |
| | | | | increase in antibiotic |
| | | | | tolerance. |
| | | | | In non-inflammatory |
| | | | | lesions, the growth of nine |
| | | | | bacterial species was |
| | | | | observed from 40 samples. |
| | Indonesia | Cross sectional study | 40 patients | In an anaerobic culture, |
| | | | | Cutibacterium acnes |
| | | | | (17,5%) was identified. In |
| | | | | aerobic cultures, different |
| Jusuf et al. ¹³ | | | | bacterial species were |
| (2020) | | | | found including |
| | | | | Staphylococcus epidermis |
| | | | | (52.5%), Staphylococcus |
| | | | | hominis (12.5%), |
| | | | | Staphylococcus |
| | | | | haemolyticus (7.5%), |
| | | | | Micrococcus luteus |
| | | | | (7.5%), Leuconostoc |
| | | | | mesentroides (7.5%), |
| | | | | 110001111 01000 (1.070); |

| | | | | Staphylococcus aureus |
|--------------------------|-------|-----------|----------|-----------------------------|
| | | | | (5%), Kocuria varians |
| | | | | (5%), and Staphylococcus |
| | | | | vitulinus (2.5%). In |
| | | | | inflammatory lesions, nine |
| | | | | bacterial species were |
| | | | | found, in which was the |
| | | | | anaerobic culture we |
| | | | | identified Cutibacterium |
| | | | | acnes (25.0%). Aerobic |
| | | | | cultures have revealed the |
| | | | | growth colonies of |
| | | | | Staphylococcus |
| | | | | epidermidis (42.5%), |
| | | | | Staphylococcus hominis |
| | | | | (22.5%), Staphylococcus |
| | | | | aureus (12.5%), |
| | | | | Staphylococcus |
| | | | | haemolyticus (10.0%), |
| | | | | Leuconostoc mesentroides |
| | | | | (5.0%), Staphylococcus |
| | | | | cohnii (2.5%), |
| | | | | Staphylococcus arlettae |
| | | | | (2.5%), and Dermacoccus |
| | | | | nishinomyaensis (2.5%). |
| | | | | Two mixed bacterial |
| | | | | growths were observed in |
| | | | | non-inflammatory lesions, |
| | | | | while four mixed bacterial |
| | | | | growths were found in |
| | | | | inflammatory lesions. |
| | | | | Targeted amplicon |
| | | | | sequencing of the 16S |
| | | | | ribosomal RNA gene |
| | Italy | Cross | 40 | (16S-seq), are enabling the |
| | | | | identification and |
| To the second | | | | quantification of |
| Finotello et | | sectional | 48 | human-resident |
| al. ¹⁴ (2018) | | study | patients | microorganisms at |
| | | | | unprecedented resolution, |
| | | | | providing novel insights |
| | | | | into the role of the |
| | | | | microbiota in health and |
| | | | | disease. Once microbial |
| | | | | |

| | | | | alarra dan aga a a a a a a a a a a a a a a a a a |
|---|--------|-----------------------------|-------------|---|
| | | | | abundances are quantified |
| | | | | through NGS data |
| | | | | analysis, diversity indices |
| | | | | provide valuable |
| | | | | mathematical tools to |
| | | | | describe the ecological |
| | | | | complexity of a single |
| | | | | • • |
| | | | | |
| | | | | _ |
| | | | | * |
| | | | | ^ * |
| | | | | |
| | | | | |
| | | | | |
| | | | | · · |
| | | | | |
| | | | | currently available. |
| | | | | 1 |
| | France | Cross sectional study | 48 patients | 1 |
| | | | | several skin dermatoses |
| | | | | physiopathology. Here, we |
| | | | | show that inflammatory |
| | | | | skin is associated with |
| | | | | changes in the skin |
| | | | | microbiota composition on |
| | | | | the back of severe acne |
| | | | | patients but also on the |
| | | | | face of patients where acne |
| Degnolie et | | | | was scored as mild to |
| 6 | | | | moderate, comparing with |
| ai. (2019) | | | | healthy controls. Changes |
| | | | | were observed particularly |
| | | | | on skin commensals |
| | | | | Propionibacteriaceae, |
| | | | | Staphylococcaceae and |
| | | | | Enterococcaceae families, |
| | | | | suggesting the importance |
| | | | | of the balance between |
| | | | | skin commensals to |
| | | | | maintain skin homeostasis |
| | | | | and control skin |
| | | | | inflammatory process. |
| Dagnelie et al. ¹⁵ (2019) | France | sectional | | sample or to detect species differences between samples. However, diversity is not a determined physical quantity for which a consensus definition and unit of measure have been established, and several diversity indices are currently available. Skin microbiota appears as a key player involved in several skin dermatoses physiopathology. Here, we show that inflammatory skin is associated with changes in the skin microbiota composition on the back of severe acne patients but also on the face of patients where acne was scored as mild to moderate, comparing with healthy controls. Changes were observed particularly on skin commensals Propionibacteriaceae, Staphylococcaceae and Enterococcaceae families, suggesting the importance of the balance between skin commensals to maintain skin homeostasis and control skin |

DISCUSSION

The discussion should be an interpretation of the results, not a repetition of the results. This discussion includes at least: an explanation of the meaning of the findings and why the findings are important; Support the answer with the results. Explain how your results relate to expectations and the literature; state clearly why the results are acceptable and whether there is any agreement or conflict with previous research results; consider alternative explanations for the findings; consider research implications; study limitations; and provide suggestions for further research.

Avoid writing in the form of bullet numbering or item list style; it is best to write it in the form of a descriptive paragraph, even though it is a list item. If it contains tables and figures, the numbering is a continuation of the previous number. Each table and figure must be given a title.

CONCLUSION

The conclusion must contain confirmation of the problems that have been analyzed in the results and discussion sections. Write a conclusion concisely and clearly. It is not recommended that the conclusion be written in several parts or points. The conclusion is intended to help readers understand why your research is important to them after they have finished reading the manuscript. A conclusion is not simply a summary of the main topics discussed or a restatement of your research problem, but rather a synthesis of the important points. It is important that the conclusion does not leave any questions unanswered.

DISCLOSURE STATEMENT

Disclosure Statement: The authors have no conflicts of Interest to declare.

REFERENCES

References should be listed in the order of their appearance in the text. Each cited source must include the author's name, article title, journal name, year of publication, volume, issue number, page numbers, and DOI (if available). Example of reference format:

1. Chen H, Zhang TC, Yin XL, Man JY, Yang XR, Lu M. Magnitude and temporal trend of acne vulgaris burden in 204 countries and territories from 1990 to 2019: an analysis from the Global Burden of Disease Study 2019. Br J Dermatol. 2022 Apr;186(4):673–83.

- 2. GBD 2019 Demographics Collaborators. Global age-sex-specific fertility, mortality, healthy life expectancy (HALE), and population estimates in 204 countries and territories, 1950-2019: a comprehensive demographic analysis for the Global Burden of Disease Study 2019. Lancet. 2020 Oct 17;396(10258):1160–203.
- 3. Walters KE, Martiny JBH. Alpha-, beta-, and gamma-diversity of bacteria varies across habitats. PLoS One. 2020;15(9):e0233872.
- 4. Dréno B, Dagnelie MA, Khammari A, Corvec S. The Skin Microbiome: A New Actor in Inflammatory Acne. Am J Clin Dermatol. 2020 Sep;21(Suppl 1):18–24.
- 5. Barnard E, Shi B, Kang D, Craft N, Li H. The balance of metagenomic elements shapes the skin microbiome in acne and health. Sci Rep. 2016 Dec 21;6:39491.