

AGES 12+ YEARS: STEPWISE APPROACH FOR MANAGEMENT OF ASTHMA

Intermittent Asthma		Management of Persistent Asthma in Individuals Ages 12+ Years				
Treatment	STEP 1	STEP 2	STEP 3	STEP 4	STEP 5	STEP 6 [■]
Preferred	PRN SABA	Daily low-dose ICS and PRN SABA or PRN concomitant ICS and SABA [▲]	Daily and PRN combination low-dose ICS-formoterol [▲]	Daily and PRN combination medium-dose ICS-formoterol [▲]	Daily medium-high dose ICS-LABA + LAMA and PRN SABA [▲]	Daily high-dose ICS-LABA + oral systemic corticosteroids + PRN SABA
Alternative		Daily LTRA [*] and PRN SABA or Cromolyn, [*] or Nedocromil, [*] or Zileuton, [*] or Theophylline, [*] and PRN SABA	Daily medium-dose ICS and PRN SABA or Daily low-dose ICS-LABA, or daily low-dose ICS + LAMA, [▲] or daily low-dose ICS + LTRA, [*] and PRN SABA or Daily low-dose ICS + Theophylline [*] or Zileuton, [*] and PRN SABA	Daily medium-dose ICS-LABA or daily medium-dose ICS + LAMA, and PRN SABA [▲] or Daily medium-dose ICS + LTRA, [*] or daily medium-dose ICS + Theophylline, [*] or daily medium-dose ICS + Zileuton, [*] and PRN SABA	Daily medium-high dose ICS-LABA or daily high-dose ICS + LTRA, [*] and PRN SABA	
		Steps 2-4: Conditionally recommend the use of subcutaneous immunotherapy as an adjunct treatment to standard pharmacotherapy in individuals ≥ 5 years of age whose asthma is controlled at the initiation, build up, and maintenance phases of immunotherapy [▲]			Consider adding Asthma Biologics (e.g., anti-IgE, anti-IL5, anti-IL5R, anti-IL4/IL13)**	

Assess Control

- First check adherence, inhaler technique, environmental factors,[▲] and comorbid conditions.
- **Step up** if needed; reassess in 2-6 weeks
- **Step down** if possible (if asthma is well controlled for at least 3 consecutive months)

Consult with asthma specialist if Step 4 or higher is required. Consider consultation at Step 3.

Control assessment is a key element of asthma care. This involves both impairment and risk. Use of objective measures, self-reported control, and health care utilization are complementary and should be employed on an ongoing basis, depending on the individual's clinical situation.

Abbreviations: ICS, inhaled corticosteroid; LABA, long-acting beta₂-agonist; LAMA, long-acting muscarinic antagonist; LTRA, leukotriene receptor antagonist; SABA, inhaled short-acting beta₂-agonist

[▲] Updated based on the 2020 guidelines.

^{*} Cromolyn, Nedocromil, LTRAs including Zileuton and montelukast, and Theophylline were not considered for this update, and/or have limited availability for use in the United States, and/or have an increased risk of adverse consequences and need for monitoring that make their use less desirable. The FDA issued a Boxed Warning for montelukast in March 2020.

^{**} The AHRQ systematic reviews that informed this report did not include studies that examined the role of asthma biologics (e.g., anti-IgE, anti-IL5, anti-IL5R, anti-IL4/IL13). Thus, this report does not contain specific recommendations for the use of biologics in asthma in Steps 5 and 6.

[■] Data on the use of LAMA therapy in individuals with severe persistent asthma (Step 6) were not included in the AHRQ systematic review and thus no recommendation is made.