

The University of Arkansas for Medical Sciences
Trauma Clinical Practice Management Guideline

SUBJECT: Pregnant Trauma Patients

AUTHORS: Kyle Kalkwarf, MD; Adam Sandlin, MD

APPROVED: 10/10/2016

UPDATED: 8/2023, 11/2025

EFFECTIVE: 11/27/2023

PURPOSE: Guidance for evaluating and treating traumatically injured pregnant patients.

DEFINITIONS:

- Fetal viability: known ≥ 22 week fetus or estimated ≥ 20 weeks
- Perimortem Caesarian delivery: surgical delivery of a fetus during cardiac arrest

GUIDELINES:

Pregnant patients with traumatic injuries, suspected of reaching a gestational age consistent with fetal viability, will be a Level 1 or 2 Trauma Activation. Additionally, emergency obstetric teams (and neonatal, if deemed necessary) will respond to the trauma bay along with the Trauma Team for these patients.

- Trauma Activations will follow the UAMS Trauma Activation Criteria Guideline. Pregnancy raises the activation level (e.g., a Level 3 activation becomes a Level 2 activation, and a Level 2 activation becomes a Level 1 activation) if the patient is pregnant with a viable fetus.
- A viable fetus is defined by **fetal viability** to accommodate for prehospital or transferring hospital underestimation of gestational age.

When a trauma patient requiring emergency obstetric response is reported en route or arrives unannounced at the trauma center, the ED Charge Nurse or designee is responsible for notifying the Emergency Obstetric, Neonatal, and Trauma Teams.

- The primary alerting mechanism activates:
 - Trauma Team group pager
 - Stork group pager
- In the event of a primary alerting mechanism failure, the emergency obstetric team can be contacted by calling Labor and Delivery at 501-526-0564, and the infant emergency group can contact the neonatal emergency team via Vocera. The OB chief resident can also be notified by calling the OB chief's cell at **501-231-6643**.

It is the intent that the evaluation of the fetus by the emergency obstetric team shall occur simultaneously with the assessment and initial management of the trauma patient by trauma and emergency medicine personnel.

- This often requires close coordination among anesthesia, emergency medicine, obstetrics, and trauma surgery attendings, as well as resident physicians.

If fetal distress or other findings warrant emergency Caesarian section, the case will be done in the

main OR trauma room (usually OR 13, but it may be a backup room). (see “TRAUMA STAT Operating Room Access”)

- Main OR rooms are equipped with all necessary Cesarean section and emergency neonatology equipment and neonatology.
- Performing the Cesarean section in the main operating room (OR) enables optimal management of other potential injuries that may arise.
- In unusual circumstances, selected Cesarean sections in Trauma patients may be performed in L&D with the concurrence of the obstetrics, trauma, and anesthesiology attendings.

A Perimortem cesarean delivery section will take place in the location where the arrest occurs because outcomes are worse if the patient is moved. This will most often be the ED trauma bay.

- The indications for perimortem c-section are:
 - Imminent maternal death
 - Immediately at the time of loss of vital signs
 - Within 4 minutes of unsuccessful CPR
- If present, the OB service will perform the Perimortem cesarean delivery with the assistance of the trauma team. It will be performed through a vertical midline laparotomy.
- Perimortem Caesarian delivery should occur before a resuscitative thoracotomy and open-heart massage because it results in better outcomes by removing the enlarged uterus, which stops the shunting of blood and improves the efficacy of CPR, which should resume after the delivery of the baby.

Trauma patients requiring admission who are pregnant at or beyond the gestational age of viability will be jointly managed by the Trauma and Obstetric services. The specific unit of admission will be determined by the services based on these guidelines:

- Patients requiring ICU care will be admitted to the SICU with continuous fetal monitoring as deemed necessary by the obstetric service.
- Patients with significant injuries but not requiring ICU admission will be admitted to the appropriate trauma unit with fetal monitoring as deemed necessary by the obstetric service.
- Patients with minor injuries will be admitted to the unit deemed appropriate by the obstetrics service, and the trauma service will follow until discharge.

Trauma patients requiring admission who are pregnant before the gestational age of viability will be managed by the trauma team with obstetric or gynecologic consultation. The specific obstetric or gynecologic care for these patients will be determined on an individual basis.

- The primary goal for these patients is to provide optimal care for the mother and child.
- The mother’s medical needs will determine where the patient is admitted. Rhogam will be considered on a case-by-case basis after consultation with the obstetric attending physician.

This guideline was prepared by the UAMS ACS Division based on a recent literature review. It is intended to support clinical decision-making and should be used at the discretion of the trauma team, as it is not a fixed policy or protocol.

REFERENCES:

1. Appelbaum RD, Yorkgitis B, Rosen J, Butts CA, To J, Knight AW, Zhang J, Kirsch JM, Levin JH, Riera KM, Kelley KM. Trauma in pregnancy: A systematic review, meta-analysis, and practice management guideline from the Eastern Association for the Surgery of Trauma. *Journal of Trauma and Acute Care Surgery*. 2025 Aug 1;99(2):298-309.
2. Liggett MR, Amro A, Son M, Schwulst S. Management of the pregnant trauma patient: a systematic literature review. *Journal of Surgical Research*. 2023 May 1;285:187-96.
3. Mendez-Figueroa H, Dahlke JD, Vrees RA, Rouse DJ. Trauma in pregnancy: an updated systematic review. *American journal of obstetrics and gynecology*. 2013 Jul 1;209(1):1-0.
4. Oxford CM, Ludmir J. Trauma in pregnancy. *Clinical obstetrics and gynecology*. 2009 Dec 1;52(4):611-29.
5. Brown HL. Trauma in pregnancy. *Obstetrics & Gynecology*. 2009 Jul 1;114(1):147-60.
6. Huls CK, Detlefs C. Trauma in pregnancy. In *Seminars in perinatology* 2018 Feb 1 (Vol. 42, No. 1, pp. 13-20). WB Saunders.
7. Mattox KL, Goetzl L. Trauma in pregnancy. *Critical care medicine*. 2005 Oct 1;33(10):S385-9.
8. Barraco RD, Chiu WC, Clancy TV, Como JJ, Ebert JB, Hess LW, Hoff WS, Holevar MR, Quirk JG, Simon BJ, Weiss PM. Practice management guidelines for the diagnosis and management of injury in the pregnant patient: the EAST Practice Management Guidelines Work Group. *Journal of Trauma and Acute Care Surgery*. 2010 Jul 1;69(1):211-4.