

Foundations I:

Trauma 2 Teaching Points

Lefort Fracture

• Case Teaching Points

The differential for facial trauma can be broken down by considering injuries to each head, face, and neck structure. This should include, but is not limited to, skull fracture, intracranial hemorrhage, orbital wall fracture, globe rupture, retrobulbar hematoma/compartment syndrome, entrapment, Le Fort fractures, mandible fracture, zygomatic arch fracture, nasal bone fracture, nasal septal hematoma, dental fracture/avulsion/subluxation, other facial fractures, auricular hematoma, laceration, foreign body, vascular injury (carotid), nerve injury, and c-spine injury

• Initial airway management in facial trauma

- Airway management in facial trauma can be challenging and requires extreme care.
- Early intubation for airway protection may be required in these cases due to extensive hemorrhage and increased risk of mechanical obstruction.
- Typical maneuvers such as jaw thrust, suctioning, or manually grasping the tongue may not be successful. Use of a nasal airway or trumpet should be avoided.
- o BVM can sometimes help stabilize the midface as it acts as a split for facial fractures

• Considerations when performing intubation

- Awake intubation should be considered first in patients with facial trauma
 - There are many ways to perform awake intubation. Importance lies in good oropharyngeal local anesthetic and appropriate anxiolysis/sedation.
- RSI is not technically contraindicated in facial trauma but should be used selectively
 given concern for difficulty with ventilation. Predicating when ventilation will be difficult
 is not always straightforward and losing the ability to ventilate can occur quickly.
 - If performing RSI, have neck prepped and cricothyrotomy supplies ready.
- You should avoid nasal intubation or blind ETT passage given distortion of typical anatomy.
- Ensure that full cervical spine precautions are used in the setting of trauma, including maintaining in-line stabilization.

• What are the different types of Le Fort fractures?

- Le Fort fractures are transverse fractures of the midface. They are divided into three classifications (sometimes four) (<u>Figure 25.2</u>).
 - Le Fort I

Revised: Fall 2022

- Transverse fracture through maxilla and pterygoid plates just below floor of the nose essentially separating teeth from upper face
- Involves alveolar ridge, lateral nose, and inferior wall of maxillary sinus
- Only hard palate and teeth move, not the nose or eyes

■ Le Fort II

- Pyramidal fracture through central maxilla and hard palate
- Involves alveolar ridge, pterygoid plates, lateral walls of maxillary sinus, inferior orbital rim, and nasal bones
- Hard palate and nose move, not the eyes

■ Le Fort III

- Craniofacial disjunction with separation of maxilla from skull base
- Involves nasofrontal suture, medial orbital wall, zygomaticofrontal suture, zygomatic arch, and pterygoid plates
- Entire face moves with exception of eyes which are held in place by optic nerve

■ Le Fort IV

- Le Fort III with frontal bone
- All require facial trauma consultation. Most require admission with IV antibiotics and potential surgical repair.
- Patients with Le Fort fractures should receive antibiotics focusing coverage on strep and anaerobes. Appropriate IV choices include clindamycin or Unasyn.

• Examination considerations and pearls in facial trauma

- A systematic examination should take place for all facial trauma so injuries are not missed.
- A few special considerations by site
 - Head/Face
 - Frontal bone fractures carry high risk of other intracranial, c-spine, or facial fracture. Consider CSF leak.
 - Isolated zygomatic fractures are rare but require surgical consult
 - Always look for multiple mandibular fractures if one is present (think of a ring). Considered open fractures until through intraoral exam. May have malocclusion. Assess for sublingual hematoma or alveolar ridge fracture with missing teeth. May cause airway compromise.

Ear

- Auricular hematoma predisposes to "cauliflower ear" and should be drained if acute.
- Assess for hemotympanum, battle sign, CSF leak.

Eyes

Blow-out fracture involves inferior or medial orbital wall. Adipose tissue, inferior rectus, or inferior oblique can become entrapped within the maxillary or ethmoid sinus. May see diplopia, inability to look up, anesthesia of cheek and upper lip, or enophthalmos.

- Lateral, inferior, or superior orbital fractures are often associated with other facial fractures.
- Naso-orbito-ethmoid fractures are often associated with ductal injuries or other traumatic brain injury.
- Consider corneal abrasion, globe rupture, orbital compartment syndrome, or retinal trauma.

Nose

- Epistaxis can be controlled with direct pressure or anterior packing.
 Caution with posterior packing as typical anatomy may be disrupted.
- Septal hematoma requires drainage given risk for septal necrosis.

Oral

- Lip lacerations require delicate repair to maintain cosmesis and function.
 May require multilayer repair.
- Through and through wounds should receive multilayer closure starting inside and work outwards.
- Ducts should be assessed for integrity prior to closure. If involved, consult facial trauma surgery.
- Tongue lacerations may only need repair if gaping, large (> 1 cm), involve distal aspect, or have significant bleeding.
- Dental injuries are common with facial trauma. Careful exam for fracture, subluxation, or avulsion should be performed in all patients.
 May require replacement and splinting. Obtain chest x-ray if teeth missing.

Neck

- C-spine precautions should be maintained until c-spine injury ruled-out
- CTA may be necessary to assess for vascular injury

Attributions

Author: Dr. Conor Dass

Editor(s): Dr. Miriam Cordeiro Stone

o Editor-in-Chief: Dr. Dana Loke, Dr. Kristen Grabow Moore

References:

- Tarlan Hedayati, Dhara P. Amin. Chapter 259: Trauma to the Face. In: Judith E. Tintinalli, O. John Ma, et al, editors. Tintinalli's Emergency Medicine: A Comprehensive Study Guide (9th ed). New York: McGraw-Hill; 2020.
- Ryanne J. Mayersak. Chapter 42: Facial Trauma. In: Ron Walls, Robert Hockberger, Marianne Gausche-Hill et al, editors. Rosen's Emergency Medicine: Concepts and Clinical Practice (9th ed). Philadelphia: Elsevier, Inc; 2018.
- Bhupendra C. Patel, Thomas Wright, Muhammad Waseem. Le Fort Fractures [Internet]. StatPearls. StatPearls Publishing; 2021 [cited 2021 Oct 15]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK526060/
- Gerry Funk. Facial Fracture Management Handbook LeFort Fractures [Internet]. Iowa Head and Neck Protocols. The Department of Otolaryngology, University of Iowa Hospitals and Clinics; 2019 [cited 2021Oct15]. Available from:

- https://medicine.uiowa.edu/iowaprotocols/facial-fracture-management-handbook-lefort-fractures
- Frank Gaillard, Dalia Ibrahim. Le Fort fracture classification. In: Radiopaedia.org [Internet]. Radiopaedia.org; 2008 [cited 2021 Nov 7]. Available from: http://radiopaedia.org/articles/1317
- Trevor Mills, Peter Deblieux. Emergency airway management in the adult with direct airway trauma. In: Post T, editor. UpToDate. [Internet]. Waltham, Mass.: UpToDate. Available from: www.uptodate.com (Accessed October 15, 2021)
- Image References
 - Maxillofacial CT from "radiopedia.org."
 https://radiopaedia.org/cases/facial-fractures-le-fort-type-ii?lang=us
 - Le Fort Fractures (figure 25.2) from https://commons.wikimedia.org/wiki/Category:Fractures_of_the_huma n_maxilla

Figure 25.2 – Le Fort Fractures

Type I	Type II	Type III
Transverse fracture through maxilla and pterygoid plates just below floor of the nose separating teeth from upper face	Pyramidal fracture through central maxilla and hard palate	Craniofacial dysjunction with separation of maxilla from skull base
Hard palate and teeth move	Maxilla and nose move	Entire face moves with exception of eyes move

Back to case teaching points

Subdural Hematoma

Case Teaching Points

The differential for AMS in an elderly patient is very broad. For this case, it should include hypoglycemia/hyperglycemia, electrolyte abnormality, hepatic encephalopathy, endocrine abnormality, hypertensive encephalopathy, overdose/polypharmacy, medication side effect, Wernicke encephalopathy, pneumonia, UTI, sepsis, stroke, intracerebral hemorrhage, CNS infection, CNS lesion, or dementia. A focused history and physical exam will help narrow down this appropriately broad initial differential diagnosis.

What is the pathophysiology of SDH, what are common clinical features, and how do you classify them?

- Pathophysiology
 - SDH is a type of intracranial hemorrhage where bleeding forms in the space between the dura and arachnoid (bone dura arachnoid pia brain).
 - Occurs due to tearing of bridging veins located in this space. These veins drain from the surface of the brain into the dural sinuses.
 - Commonly occurs in the elderly and alcoholic patients as the vessels travel further distances given atrophy.
- Common clinical features
 - Clinical presentation has a wide spectrum ranging from completely asymptomatic to comatose. Symptoms may not develop until several days to weeks after the initial injury as venous bleeding is slow. Symptoms can be from initial injury or from ischemia/damage caused by prolonged compression of brain structures from the slowly expanding hematoma or from midline shift.
 - Acute SDH may present with a decreased level of consciousness. Common symptoms for subacute and chronic SDH include headache, photophobia, altered mental status, weakness, or other vague symptoms.
- Classification

Acute: < 2 daysSubacute: 3-14 daysChronic: > 15 days

What do you expect to see on head CT?

- Depends on the timing of the injury (Figure 38.5).
- Subacute injury is often the most difficult to identify as the collection becomes isodense
 with surrounding structures. Secondary signs such as sulcal effacement or mass effect
 are helpful clues to consider. Still, CT with contrast or MRI may be necessary for further
 characterization.
- You may also see acute-on-chronic findings within the SDH.

• How do you manage SDH in the emergency department?

- Initial management should focus on stabilization including ABCs and quick neurosurgical consultation.
- Other important early considerations include:
 - Avoiding hypotension <u>or</u> hypertension.
 - MAP goal > 75-80 and the SBP goal is generally 100-110 to 140-160 (110 if age 15-49 or ≥70 years). This may require IV fluids, vasopressors, or vasodilators to maintain.
 - HOB 30 degrees
 - Reverse anticoagulation
 - Consider platelet transfusion only if plts < 100,000
 - Avoid hypoxia
 - Avoid hyperthermia
 - Avoid hypoglycemia
 - Consider antiepileptics
- Surgical intervention is recommended in patients with:
 - SDH width > 10 mm,
 - Midline shift > 5 mm, or
 - GCS < 9 or change ≥ 2 since injury</p>
- Temporizing measures are discussed further below

What are the common herniation syndromes (<u>Figure 38.6</u>) and what are the treatment options?

- Subfalcine herniation
 - Presents similar to ACA stroke with contralateral leg weakness as ACA gets compressed
- Central transtentorial herniation
 - Pupils initially miotic but progress to fixed and misposition
 - Decorticate and then decerebrate posturing
 - Pathologic breathing patterns (ex. Cheyne-Stokes)
- Uncal herniation
 - Loss of consciousness (ascending arousal pathway), ipsilateral pupillary dilation (oculomotor nerve), and contralateral hemiparesis (corticospinal tract). May have positive Babinski.
 - May have false localizing symptoms and can progress to include bilateral symptoms and posturing
- Cerebellar tonsillar herniation
 - Miotic and unreactive pupils, posturing to flaccid paralysis, and cardiovascular/respiratory collapse
- Treatment options
 - HOB 30 degrees
 - Osmotherapy

- Hypertonic saline Bolus 250 mL of 3%, can be given through peripheral line
- Mannitol (may be inferior to hypertonic saline) − 1 g/kg bolus
- Hyperventilation with target pCO2 35-40
- Emergent neurosurgical intervention

What are the other types of intracranial hemorrhage?

- o Epidural hematoma
- Subarachnoid hemorrhage
- Intraparenchymal hemorrhage

POCUS Pearls (time permitting)

- Ocular US has many uses in the ED. Optic nerve sheath diameter (ONSD) can be used to assess for evidence of increased intracranial pressure. The optic nerve sheath is contiguous with the subarachnoid space, therefore it distends with increased intracranial pressure.
- Fundoscopy can be very difficult for EMPs. Measuring the ONSD is an easier and more reliable way of assessing for increased ICP.
- The optic nerve appears hypoechoic and the sheath appears hyperechoic; ONSD should be measured 3 mm posterior to the globe, from outer edge to outer edge
- In adults, a diameter > 5 mm is abnormal and consistent with increased ICP, though some data suggest a cutoff of 6mm.
- See <u>Figure 38.4</u> for example of abnormal Optic Nerve Sheath Diameter (ONSD) measurement, consistent with increased ICP.

Attributions

• Author: Dr. Conor Dass

o Editor(s): Dr. Miriam Cordeiro Stone

o Editor-in-Chief: Dr. Dana Loke, Dr. Kristen Grabow Moore

• Ultrasound content by: Dr. Rachel Haney, Dr. Sierra Beck

References:

- David W. Wright, Lisa H. Merck. Chapter 257: Head Trauma. In: Judith E. Tintinalli, O. John Ma, et al, editors. Tintinalli's Emergency Medicine: A Comprehensive Study Guide (9th ed). New York: McGraw-Hill; 2020.
- William G. Heegaard, Michelle H. Biros. Chapter 41: Head Injury. In: Ron Walls, Robert Hockberger, Marianne Gausche-Hill et al, editors. Rosen's Emergency Medicine: Concepts and Clinical Practice (9th ed). Philadelphia: Elsevier, Inc; 2018.
- William McBride. Subdural hematoma in adults: Etiology, clinical features, and diagnosis. In: Post T, editor. UpToDate. [Internet]. Waltham, Mass.: UpToDate. Available from: www.uptodate.com (Accessed October 6, 2021)
- William McBride. Subdural hematoma in adults: prognosis and management. In: Post T, editor. UpToDate. [Internet]. Waltham, Mass.: UpToDate. Available from: www.uptodate.com (Accessed October 6, 2021)

- William Shy, Debbie Yi Madhok. Update on the ED management of intracranial hemorrhage: Not all head bleeds are the same [Internet]. Update on the ED Management of Intracranial Hemorrhage: Not All Head Bleeds Are the Same. ALIEM; 2017 [cited 2021Oct15]. Available from: https://www.aliem.com/intracranial-hemorrhage-management/
- Farkas J. Approach to stupor & amp; coma [Internet]. Approach to Stupor & amp; Coma. The Internet Book of Critical Care; 2021 [cited 2021Oct15]. Available from: https://emcrit.org/ibcc/coma/#common_coma_syndromes
- Sunil Munakomi, Joe M Das. Brain Herniation. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK542246/
- Image References
 - Figure 38.1: ECG from "Life in the Fastlane" https://litfl.com/normal-sinus-rhythm-ecg-library/
 - Figure 38.2: Chest Xray from "radiopedia" https://radiopaedia.org/cases/normal-chest-x-ray-1?lang=us
 - Figures 38.3, 38.5: Head CT's with descriptions, from "radiopedia" https://radiopaedia.org/cases/subdural-haemorrhage-6?lang=us, https://radiopaedia.org/cases/subacute-subdural-haematoma, https://radiopaedia.org/cases/chronic-subdural-haemorrhage-teaching? lang=us
 - Figure 38.4 Ocular POCUS images courtesy of: Northwestern Emergency Medicine POCUS Image Bank AND/OR Emory Emergency Medicine POCUS Archive
 - Figure 38.6: Brain herniation syndromes from https://commons.wikimedia.org/wiki/File:Brain_herniation_types-2.svg

Figure 38.4 – POCUS Ocular US




Back to case

Figure 38.5 – SDH CT Findings by Age

Age	Image	Typical Findings
Acute (1-2 days)		Highly visible crescent-shaped hyperdense collection. May see midline shift.
Subacute (3-14 days)	R RD: 250 RD: 250 RD: 250 RD: 250 ROy: 120 ROy: 120 ROy: 252 ROy: 25	As clot degradation begins, the collection becomes hypodense making it potentially difficult to identify. Secondary signs are important to recognize, including: 1. Sulci do not reach the skull border 2. Mass-effect including sulcal effacement/distortion (gyri appear pushed together) and midline shift
Chronic (>15 days)		Hypodense crescent-shaped collection. May see midline shift.

Back to case teaching points

Figure 38.6 – Brain Herniation Syndromes



Back to case teaching points

Neurogenic Shock

• Case Teaching Points

 The patient in this case is a trauma patient presenting with unstable vital signs and neurologic compromise. An appropriate trauma differential diagnosis should be considered. In blunt trauma, spinal injuries should be presumed and should be treated as unstable until proven otherwise. In cases of low clinical suspicion, Nexus and Canadian C-spine rules are helpful in determining if imaging is needed.

• How can you evaluate clinically for suspected spinal cord injury?

- Determine sensory level (pinprick and light touch):
 - C6 thumb
 - C7 middle finger
 - C8 little finger
 - T4 nipple line
 - T10 umbilicus
 - T12 inguinal ligament
 - L2 anterior thigh
 - L4 medial malleolus
 - S1 lateral heel
 - S4/5 perianal area
- Determine motor level (against gravity):
 - C5 elbow flexion
 - C6 wrist extension
 - C7 elbow extension
 - C8 finger flexion
 - T1 5th digit abduction
 - L2 hip flexion
 - L3 knee extension
 - L4 dorsiflexion
 - L5 toe extension
 - S1 plantar flexion
- Check for sacral sparing (suggests incomplete spinal cord injury): perineal sensation, rectal tone, anal wink (touching surrounding skin causes anal contraction), bulbocavernosus reflex (squeezing glans/clitoris or tugging Foley causes anal contraction) which are all absent in spinal shock. Return of function signifies end of spinal shock period.

• What types of shock should you assess for in patients with spinal cord injuries?

 Neurogenic shock: loss of sympathetic vascular tone leading to bradycardia and hypotension from unopposed parasympathetic (vagal) tone. Occurs only in patients with spinal cord injuries above T6; if below T6 assume hemorrhagic shock is the cause of hypotension.

- Spinal "shock": spinal cord concussion (not cardiovascular in origin) leading to 24-72 hr period of paralysis, hypotonia, and areflexia. The end of this period may be indicated by the return of rectal tone. Should have stable vital signs and thus is not true shock in the classic sense.
- Hemorrhagic shock: may be difficult to assess in patients with neurogenic shock due to autonomic dysfunction and blunting of normal tachycardia. Shock in trauma is presumed to be hemorrhagic until proven otherwise.

• What is the treatment for spinal cord injuries?

- Intubate early for signs of respiratory failure if there is concern for diaphragmatic involvement (phrenic nerve injury from lesions above C5).
- Place Foley to decompress a distended, neurogenic bladder
- Steroids are controversial (initial studies showed improved neurologic outcomes if high dose steroids are given within 8 hr, while more recent studies show a substantial risk of infection and avascular necrosis). This decision should be made in consultation with your spinal surgeon.
- Treat neurogenic shock as needed.

• How do you treat neurogenic shock?

- Avoid hypoxia or hypotension to protect spinal cord perfusion.
- Fluid resuscitation should be attempted before vasopressors but caution to avoid pulmonary edema and ARDS.
 - Maintain urine output > 0.5 mL/kg/h.
- Goal MAP 85-90 mmHg, give vasopressors as needed
 - Norepinephrine is the treatment of choice
 - Phenylephrine can be added, but only after norepinephrine has been started, as phenylephrine monotherapy may improve blood pressure but will cause reflex bradycardia.
- Give atropine as needed to maintain HR > 60.
- Watch out for hypothermia as peripheral vasodilation can lead to significant heat loss.
- Thromboembolism is very common due to pooling of blood in a dilated venous system. Consider early initiation of anticoagulation once hemorrhage has been ruled out.

Attributions

Author: Dr. Andrew Mastanduono

o Editor(s): Dr. Miriam Stone

o Editor-in-Chief: Dr. Dana Loke, Dr. Kristen Grabow Moore

References:

Gatz JD, Whetstone W. EM:RAP CorePendium [Internet]. EM:RAP CorePendium. 2021 [cited 2021Aug9]. Available from:
 https://www.emrap.org/corependium/chapter/recxnd6zjQJCQ0K5Q/Spinal-Trauma

- Steven Go. Chapter 258: Spine Trauma. In: Judith E. Tintinalli, O. John Ma, et al, editors. Tintinalli's Emergency Medicine: A Comprehensive Study Guide (9th ed). New York: McGraw-Hill; 2020.
- David M. Somand; Kevin R. Ward. Chapter 13: Approach to Traumatic Shock. In: Judith E. Tintinalli, O. John Ma, et al, editors. Tintinalli's Emergency Medicine: A Comprehensive Study Guide (9th ed). New York: McGraw-Hill; 2020.
- CT scan from Radiopedia: https://radiopaedia.org/cases/cervical-trauma-with-burst-fracture-2