EXERCISE 12.1 PAGE: 234

1. Get the algebraic expressions in the following cases using variables, constants and arithmetic operations.

(i) Subtraction of z from y. Solution:-

$$= Y - z$$

(ii) One-half of the sum of numbers x and y. Solution:-

$$= \frac{1}{2} (x + y)$$
  
=  $(x + y)/2$ 

(iii) The number z multiplied by

itself. Solution:-

$$= z \times z$$
$$= z^2$$

(iv) One-fourth of the product of numbers p and q. Solution:-

$$= \frac{1}{4} (p \times q)$$
$$= pq/4$$

(v) Numbers x and y both squared and added. Solution:-

$$= x^2 + y^2$$

(vi) Number 5 added to three times the product of numbers m and n. Solution:-

$$= 3mn + 5$$

(vii) Product of numbers y and z subtracted from

10. Solution:-

$$= 10 - (y \times z)$$
$$= 10 - yz$$

(viii) Sum of numbers a and b subtracted from their product.

## Solution:-

$$= (a \times b) - (a + b)$$
  
=  $ab - (a + b)$ 


2. (i) Identify the terms and their factors in the following expressions Show the terms and factors by tree diagrams.

(a) 
$$x-3$$

# Solution:-

Expression: x - 3

Terms: x, -3 Factors: x; -3




# (b) $1 + x + x^2$

## Solution:-

Expression:  $1 + x + x^2$ 

Terms: 1, x, x<sup>2</sup> Factors: 1; x; x,x




# (c) $y-y^3$

# **Solution:-**

Expression:  $y - y^3$ 

Terms: y, -y<sup>3</sup>

Factors: y; -y, -y, -y




# (d) $5xy^2 + 7x^2y$

# Solution:-

Expression:  $5xy^2 + 7x^2y$ 

Terms:  $5xy^2$ ,  $7x^2y$ 

Factors: 5, x, y, y; 7, x, x, y




(e) 
$$-ab + 2b^2 - 3a^2$$

# **Solution:-**

Expression:  $-ab + 2b^2 - 3a^2$ 

Terms: -ab, 2b<sup>2</sup>, -3a<sup>2</sup>

Factors: -a, b; 2, b, b; -3, a, a



(ii) Identify terms and factors in the expressions given below:

$$(a) - 4x + 5$$

(b) 
$$-4x + 5y$$

(c) 
$$5y + 3y^2$$

(d) 
$$xy + 2x^2y^2$$

$$(e) pq + q$$

(f) 1.2 ab 
$$-$$
 2.4 b + 3.6 a (g)  $\frac{3}{4}$  x +  $\frac{1}{4}$ 

(h) 
$$0.1 p^2 + 0.2 q^2$$

## Solution:-

Expressions is defined as, numbers, symbols and operators (such as +. - ,  $\times$  and  $\div$ ) grouped together that show the value of something.

In algebra a term is either a single number or variable, or numbers and variables multiplied together. Terms are separated by + or – signs or sometimes by division.

Factors is defined as, numbers we can multiply together to get another number.

SI.No.	Expression	Terms	Factors
(a)	- 4x + 5	-4x	-4, x
		5	5
(b)	– 4x + 5y	-4x	-4, x
		5у	5, y
(c)	5y + 3y <sup>2</sup>	5у	5, y
		3y²	3, y, y
(d)	$xy + 2x^2y^2$	ху	х, у
		$2x^2y^2$	2, x, x, y, y

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

(e)	pq + q	pq	P, q
		q	Q
(f)	1.2 ab – 2.4 b +	1.2ab	1.2, a, b
	3.6 a	-2.4b	-2.4, b
		3.6a	3.6, a

(g)	3/4 X + 1/4	3/4	Х	³⁄4, X
		1/2	í	1/4
(h)	$0.1 p^2 + 0.2 q^2$	1.1	p <sup>2</sup>	1.1, p, p
		1.2	$q^2$	1.2, q, q

# 3. Identify the numerical coefficients of terms (other than constants) in the following expressions:

(i) 
$$5 - 3t^2$$

(ii) 
$$1 + t + t^2 + t^3$$

(iii) 
$$x + 2xy + 3y$$

$$(v) - p^2q^2 + 7pq$$

(vi) 
$$1.2 a + 0.8 b$$

(ix) 
$$0.1 y + 0.01 y^2$$

#### Solution:-

Expressions is defined as, numbers, symbols and operators (such as +. - ,  $\times$  and  $\div$ ) grouped together that show the value of something.

In algebra a term is either a single number or variable, or numbers and variables multiplied together. Terms are separated by + or – signs or sometimes by division.

A coefficient is a number used to multiply a variable (2x means 2 times x, so 2 is a coefficient) Variables on their own (without a number next to them) actually have a coefficient of 1 (x is really 1x)

Sl.No.	Expression	Terms	Coefficients
(i)	5 – 3t <sup>2</sup>	- 3t <sup>2</sup>	-3
(ii)	$1 + t + t^2 + t^3$	t	1
		t 2	1 1
		<del>†</del> 3	
(iii)	x + 2xy + 3y	х	1
		2xy	2
		Зу	3
(iv)	100m + 1000n	100m	100

NCERT Solutions for Class 7 Maths Chapter 12

At large to see 2 or	Europe and a mar
Algebraic	Expressions

		1000n	1000
(v)	$-p^2q^2 + 7pq$	-p <sup>2</sup> q <sup>2</sup>	-1
		7pq	7
(vi)	1.2 a + 0.8 b	1.2a	1.2
		0.8b	0.8
(vii)	3.14 r <sup>2</sup>	3.14 <sup>2</sup>	3.14

(viii)	2 (l + b)	21	2
		2b	2
(ix)	$0.1 \text{ y} + 0.01 \text{ y}^2$	0.1y	0.1
		$0.01y^2$	0.01

# 4. (a) Identify terms which contain x and give the coefficient of x.

(i)  $y^2x + y$ 

(ii)  $13y^2 - 8yx$ 

(iii) x + y + 2

(iv) 5 + z + zx

(v) 1 + x + xy

(vi)  $12xy^2 + 25$ 

(vii)  $7x + xy^2$ 

## Solution:-

SI.No.	Expression	Terms	Coefficient of x
(i)	$y^2x + y$	y <sup>2</sup> x	V <sup>2</sup>
(ii)	13y² – 8yx	- 8yx	-8y
(iii)	x + y + 2	х	1
(iv)	5 + z + zx	х	1
		ZX	Z
(v)	1 + x + xy	ху	У
(vi)	12xy <sup>2</sup> + 25	12xy²	12y²
(vii)	$7x + xy^2$	7x	7
		xy <sup>2</sup>	$V^2$

# (b) Identify terms which contain $y^2$ and give the coefficient of $y^2$ .

(i)  $8 - xy^2$ 

(ii)  $5y^2 + 7x$ 

(iii)  $2x^2y - 15xy^2 + 7y^2$ 

## Solution:-

Sl.No.	Expression	Terms	Coefficient of x
(i)	8 – xy²	- xy²	- x
(ii)	$5y^2 + 7x$	5y²	5
(iii)	$2x^2y - 15xy^2 + 7y^2$	– 15xy²	– 15x
		7y²	7

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

# 5. Classify into monomials, binomials and trinomials.

(i) **4y** 

– 7z

Solution:-

Binomial.

An expression which contains two unlike terms is called a binomial.

(ii	١	
	١	\ '°
		v

## **Solution:-**

Monomial.

An expression with only one term is called a monomial.

## (iii) x + y -

## xy Solution:-

Trinomial.

An expression which contains three terms is called a trinomial.

## (iv) **100**

## **Solution:-**

Monomial.

An expression with only one term is called a monomial.

$$(v)$$
 ab – a

**–** b

## Solution:-

Trinomial.

An expression which contains three terms is called a trinomial.

3t

#### Solution:-

Binomial.

An expression which contains two unlike terms is called a binomial.

# (vii) $4p^2q -$

# 4pq<sup>2</sup> Solution:-

Binomial.

An expression which contains two unlike terms is called a binomial.

## (viii) 7mn

## Solution:-

Monomial.

An expression with only one term is called a monomial.

(ix) 
$$z^2 - 3z$$

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

+8

Solution:-

Trinomial.

An expression which contains three terms is called a trinomial.

$$\begin{array}{ccc} (x) & a^2 \\ + & b^2 \end{array}$$

#### Solution:-

Binomial.

An expression which contains two unlike terms is called a binomial.

(xi) 
$$z^2 +$$

Z

## **Solution:-**

Binomial.

An expression which contains two unlike terms is called a binomial.

(xii) 
$$1 + x +$$

## x<sup>2</sup> Solution:-

Trinomial.

An expression which contains three terms is called a trinomial.

# 6. State whether a given pair of terms is of like or unlike

terms. (i) 1, 100

Solution:-

Like term.

When term have the same algebraic factors, they are like terms.

(ii) 
$$-7x$$
,  $(5/2)x$ 

#### Solution:-

Like term.

When term have the same algebraic factors, they are like terms.

(iii) 
$$-29x, -$$

## 29y Solution:-

Unlike terms.

The terms have different algebraic factors, they are unlike terms.

42yx

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

# **Solution:-**

Like term.

When term have the same algebraic factors, they are like terms.

### Solution:-

Unlike terms.

The terms have different algebraic factors, they are unlike terms.

## Solution:-

Unlike terms.

The terms have different algebraic factors, they are unlike terms.

## 7. Identify like terms in the following:

(a) 
$$-xy^2$$
,  $-4yx^2$ ,  $8x^2$ ,  $2xy^2$ ,  $7y$ ,  $-11x^2$ ,  $-100x$ ,  $-11yx$ ,  $20x^2y$ ,  $-6x^2$ ,  $y$ ,  $2xy$ ,  $3x$  Solution:-

When term have the same algebraic factors, they are like terms.

They are,

$$-xy^{2}$$
,  $2xy^{2}$   
 $-4yx^{2}$ ,  $20x^{2}y$   
 $8x^{2}$ ,  $-11x^{2}$ ,  $-6x^{2}$   
 $7y$ ,  $y$   
 $-100x$ ,  $3x$   
 $-11yx$ ,  $2xy$ 

(b) 
$$10pq$$
,  $7p$ ,  $8q$ ,  $-p^2q^2$ ,  $-7qp$ ,  $-100q$ ,  $-23$ ,  $12q^2p^2$ ,  $-5p^2$ ,  $41$ ,  $2405p$ ,  $78qp$ ,  $13p^2q$ ,  $qp^2$ ,  $701p^2$ 

#### Solution:-

When term have the same algebraic factors, they are like terms.

They are,

EXERCISE 12.2 PAGE: 239

## 1. Simplify combining like terms:

## (i) 21b - 32 + 7b - 20b

#### Solution:-

When term have the same algebraic factors, they are like terms.

Then,

(ii) 
$$-z^2 + 13z^2 - 5z + 7z^3 - 15z$$

#### Solution:-

When term have the same algebraic factors, they are like terms.

Then,

$$= 7z3 + (-z2 + 13z2) + (-5z - 15z)$$

$$= 7z3 + z2 (-1 + 13) + z (-5 - 15)$$

$$= 7z3 + z2 (12) + z (-20) + 7z3$$

$$= 7z3 + 12z2 - 20z + 7z3$$

(iii) 
$$p - (p - q) - q - (q - q)$$

## p) Solution:-

When term have the same algebraic factors, they are like terms.

Then,

$$= p - p + q - q - q + p$$
  
 $= p - q$ 

(iv) 
$$3a-2b-ab-(a-b+ab)+3ab+b$$

#### - a Solution:-

When term have the same algebraic factors, they are like terms.

Then,

$$= 3a - 2b - ab - a + b - ab + 3ab + b - a$$
  
 $= 3a - a - a - 2b + b + b - ab - ab + 3ab$   
 $= a (3 - 1 - 1) + b (-2 + 1 + 1) + ab (-1 - 1 + 3)$   
 $= a (3 - 2) + b (-2 + 2) + ab (-2 + 3)$ 

$$= a (1) + b (0) + ab (1)$$
  
 $= a + ab$ 

(v) 
$$5x^2y - 5x^2 + 3yx^2 - 3y^2 + x^2 - y^2 + 8xy^2 - 3y^2$$
  
Solution:-

When term have the same algebraic factors, they are like terms.

Then,

$$= 5x^{2}y + 3yx^{2} - 5x^{2} + x^{2} - 3y^{2} - y^{2} - 3y^{2}$$

$$= x^{2}y (5 + 3) + x^{2} (-5 + 1) + y^{2} (-3 - 1 - 3) + 8xy^{2}$$

$$= x^{2}y (8) + x^{2} (-4) + y^{2} (-7) + 8xy^{2}$$

$$= 8x^{2}y - 4x^{2} - 7y^{2} + 8xy^{2}$$

(vi) 
$$(3y^2 + 5y - 4) - (8y - y^2 - 4)$$

### Solution:-

When term have the same algebraic factors, they are like terms.

Then,

$$= 3y^{2} + 5y - 4 - 8y + y^{2} + 4$$

$$= 3y^{2} + y^{2} + 5y - 8y - 4 + 4$$

$$= y^{2} (3 + 1) + y (5 - 8) + (-4 + 4)$$

$$= y^{2} (4) + y (-3) + (0)$$

$$= 4y^{2} - 3y$$

### 2. Add:

# (i) 3mn, – 5mn, 8mn, –

#### 4mn Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

## (ii) t - 8tz, 3tz - z, z

#### - t Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= t - 8tz + (3tz - z) + (z - t)$$

$$= t - 8tz + 3tz - z + z - t$$

$$= t - t - 8tz + 3tz - z + z$$

$$= t (1 - 1) + tz (-8 + 3) + z (-1 + 1)$$

$$= t (0) + tz (-5) + z (0)$$

$$= -5tz$$

## (iii) – 7mn + 5, 12mn + 2, 9mn – 8, – 2mn – 3 Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

# (iv) a + b - 3, b - a + 3, a - b + 3 Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= a + b - 3 + (b - a + 3) + (a - b + 3)$$

$$= a + b - 3 + b - a + 3 + a - b + 3$$

$$= a - a + a + b + b - b - 3 + 3 + 3$$

$$= a (1 - 1 + 1) + b (1 + 1 - 1) + (-3 + 3 + 3)$$

$$= a (2 - 1) + b (2 - 1) + (-3 + 6)$$

$$= a (1) + b (1) + (3)$$

$$= a + b + 3$$

# (v) 14x + 10y - 12xy - 13, 18 - 7x - 10y + 8xy, 4xy Solution:-

When term have the same algebraic factors, they are like terms. Then, we have to add the like terms

$$= 14x + 10y - 12xy - 13 + (18 - 7x - 10y + 8xy) + 4xy$$

$$= 14x + 10y - 12xy - 13 + 18 - 7x - 10y + 8xy + 4xy$$

$$= 14x - 7x + 10y - 10y - 12xy + 8xy + 4xy - 13 + 18$$

$$= x (14 - 7) + y (10 - 10) + xy(-12 + 8 + 4) + (-13 + 18)$$

$$= x (7) + y (0) + xy(0) + (5)$$

$$= 7x + 5$$

## (vi) 5m - 7n, 3n - 4m + 2, 2m - 3mn - 5 Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

# (vii) $4x^2y$ , $-3xy^2$ , $-5xy^2$ , $5x^2y$ Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= 4x^{2}y + (-3xy^{2}) + (-5xy^{2}) + 5x^{2}y$$

$$= 4x^{2}y + 5x^{2}y - 3xy^{2} - 5xy^{2}$$

$$= x^{2}y (4 + 5) + xy^{2} (-3 - 5)$$

$$= x^{2}y (9) + xy^{2} (-8)$$

$$= 9x^{2}y - 8xy^{2}$$

(viii) 
$$3p^2q^2 - 4pq + 5$$
,  $-10 p^2q^2$ ,  $15 + 9pq + 7p^2q^2$   
Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= 3p^{2}q^{2} - 4pq + 5 + (-10p^{2}q^{2}) + 15 + 9pq + 7p^{2}q^{2}$$

$$= 3p^{2}q^{2} - 10p^{2}q^{2} + 7p^{2}q^{2} - 4pq + 9pq + 5 + 15$$

$$= p^{2}q^{2} (3 - 10 + 7) + pq (-4 + 9) + (5 + 15)$$

$$= p^{2}q^{2} (0) + pq (5) + 20$$

$$= 5pq + 20$$

### 4b Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= ab - 4a + (4b - ab) + (4a - 4b)$$

$$= ab - 4a + 4b - ab + 4a - 4b$$

$$= ab - ab - 4a + 4a + 4b - 4b$$

$$= ab (1 - 1) + a (4 - 4) + b (4 - 4)$$

$$= ab (0) + a (0) + b (0)$$

$$= 0$$

(x) 
$$x^2 - y^2 - 1$$
,  $y^2 - 1 - x^2$ ,  $1 - x^2 - y^2$ 

### Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to add the like terms

$$= x^{2} - y^{2} - 1 + (y^{2} - 1 - x^{2}) + (1 - x^{2} - y^{2})$$

$$= x^{2} - y^{2} - 1 + y^{2} - 1 - x^{2} + 1 - x^{2} - y^{2}$$

$$= x^{2} - x^{2} - x^{2} - y^{2} + y^{2} - y^{2} - 1 - 1 + 1$$

$$= x^{2} (1 - 1 - 1) + y^{2} (-1 + 1 - 1) + (-1 - 1 + 1)$$

$$= x^{2} (1 - 2) + y^{2} (-2 + 1) + (-2 + 1)$$

$$= x^{2} (-1) + y^{2} (-1) + (-1)$$

$$= -x^{2} - y^{2} - 1$$

### 3. Subtract:

(i) 
$$-5y^2$$

# from y<sup>2</sup>

#### Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

$$= y^{2} - (-5y^{2})$$
$$= y^{2} + 5y^{2}$$
$$= 6y^{2}$$

# (ii) 6xy from

## -12xy Solution:-

When term have the same algebraic factors, they are like terms. Then, we have to subtract the like terms

$$= -12xy - 6xy$$

$$= -18xy$$

# (iii) (a - b) from (a +

## b) Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

# (iv) a (b-5) from b (5-

## a) Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

# (v) $-m^2 + 5mn \text{ from } 4m^2 - 3mn$

#### +8 Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

$$= 4m^{2} - 3mn + 8 - (-m^{2} + 5mn)$$

$$= 4m^{2} - 3mn + 8 + m^{2} - 5mn$$

$$= 4m^{2} + m^{2} - 3mn - 5mn + 8$$

$$= 3m^{2} - 8mn + 8$$

(vi) 
$$-x^2 + 10x - 5$$
 from  $5x -$ 

#### 10 Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

$$= 5x - 10 - (-x^2 + 10x - 5)$$
$$= 5x - 10 + x^2 - 10x + 5$$

$$= x^2 + 5x - 10x - 10 + 5$$
$$= x^2 - 5x - 5$$

# (vii) $5a^2 - 7ab + 5b^2$ from $3ab - 2a^2 - 2b^2$ Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

$$= 3ab - 2a^{2} - 2b^{2} - (5a^{2} - 7ab + 5b^{2})$$

$$= 3ab - 2a^{2} - 2b^{2} - 5a^{2} + 7ab - 5b^{2}$$

$$= 3ab + 7ab - 2a^{2} - 5a^{2} - 2b^{2} - 5b^{2}$$

$$= 10ab - 7a^{2} - 7b^{2}$$

# (viii) $4pq - 5q^2 - 3p^2$ from $5p^2 + 3q^2 - pq$ Solution:-

When term have the same algebraic factors, they are like terms.

Then, we have to subtract the like terms

$$= 5p^{2} + 3q^{2} - pq - (4pq - 5q^{2} - 3p^{2})$$

$$= 5p^{2} + 3q^{2} - pq - 4pq + 5q^{2} + 3p^{2}$$

$$= 5p^{2} + 3p^{2} + 3q^{2} + 5q^{2} - pq - 4pq$$

$$= 8p^{2} + 8q^{2} - 5pq$$

# 4. (a) What should be added to $x^2 + xy + y^2$ to obtain $2x^2 + 3xy$ ? Solution:-

Let us assume p be the required term

Then,

$$p + (x^{2} + xy + y^{2}) = 2x^{2} + 3xy$$

$$p = (2x^{2} + 3xy) - (x^{2} + xy + y^{2})$$

$$p = 2x^{2} + 3xy - x^{2} - xy - y^{2}$$

$$p = 2x^{2} - x^{2} + 3xy - xy - y^{2}$$

$$p = x^{2} - 2xy - y^{2}$$

# (b) What should be subtracted from 2a + 8b + 10 to get -3a + 7b + 16? Solution:-

Let us assume x be the required term

Then,

$$2a + 8b + 10 - x = -3a + 7b + 16$$
  
 $x = (2a + 8b + 10) - (-3a + 7b + 16)$ 

$$x = 2a + 8b + 10 + 3a - 7b - 16$$
  
 $x = 2a + 3a + 8b - 7b + 10 - 16$   
 $x = 5a + b - 6$ 

# 5. What should be taken away from $3x^2 - 4y^2 + 5xy + 20$ to obtain $-x^2 - y^2 + 6xy + 20$ ? Solution:-

Let us assume a be the required term

Then,

$$3x^{2} - 4y^{2} + 5xy + 20 - a = -x^{2} - y^{2} + 6xy + 20$$

$$a = 3x^{2} - 4y^{2} + 5xy + 20 - (-x^{2} - y^{2} + 6xy + 20)$$

$$a = 3x^{2} - 4y^{2} + 5xy + 20 + x^{2} + y^{2} - 6xy - 20$$

$$a = 3x^{2} + x^{2} - 4y^{2} + y^{2} + 5xy - 6xy + 20 - 20$$

$$a = 4x^{2} - 3y^{2} - xy$$

# 6. (a) From the sum of 3x - y + 11 and -y - 11, subtract 3x - y - 11. Solution:-

First we have to find out the sum of 3x - y + 11 and -y - 11

$$= 3x - y + 11 + (-y - 11)$$

$$= 3x - y + 11 - y - 11$$

$$= 3x - y - y + 11 - 11$$

$$= 3x - 2y$$

Now, subtract 3x - y - 11 from 3x - 2y

$$= 3x - 2y - (3x - y - 11)$$
  
= 3x - 2y - 3x + y + 11  
= 3x - 3x - 2y + y + 11

$$= -y + 11$$

# (b) From the sum of 4 + 3x and $5 - 4x + 2x^2$ , subtract the sum of $3x^2 - 5x$ and $-x^2 + 2x + 5$ .

Solution:-

First we have to find out the sum of 4 + 3x and  $5 - 4x + 2x^2$ 

$$= 4 + 3x + (5 - 4x + 2x^{2})$$

$$= 4 + 3x + 5 - 4x + 2x^{2}$$

$$= 4 + 5 + 3x - 4x + 2x^{2}$$

$$= 9 - x + 2x^{2}$$

$$= 2x^{2} - x + 9 \qquad \dots \text{ [equation 1]}$$

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

Then, we have to find out the sum of  $3x^2 - 5x$  and  $-x^2 + 2x + 5$ 

$$= 3x^{2} - 5x + (-x^{2} + 2x + 5)$$

$$= 3x^{2} - 5x - x^{2} + 2x + 5$$

$$= 3x^{2} - x^{2} - 5x + 2x + 5$$

$$= 2x^{2} - 3x + 5 \qquad ... [equation 2]$$
Now, we have to subtract equation (2) from equation (1)
$$= 2x^{2} - x + 9 - (2x^{2} - 3x + 5)$$

$$= 2x^{2} - x + 9 - 2x^{2} + 3x - 5$$

$$= 2x^{2} - 2x^{2} - x + 3x + 9 - 5$$

$$= 2x + 4$$

EXERCISE 12.3 PAGE: 242

## 1. If m = 2, find the value of:

**-2** 

### Solution:-

From the question it is given that m = 2

Then, substitute the value of m in the question

= 0

## (ii) 3m

**-** 5

#### Solution:-

From the question it is given that m = 2

Then, substitute the value of m in the question

$$= (3 \times 2) - 5$$

$$= 6 - 5$$

= 1

## (iii) 9 –

5m

#### Solution:-

From the question it is given that m = 2

Then, substitute the value of m in the question

$$= 9 - (5 \times 2)$$

$$= 9 - 10$$

# (iv) $3m^2 - 2m$

## - 7 Solution:-

From the question it is given that m = 2

Then, substitute the value of m in the question

$$= (3 \times 2^2) - (2 \times 2) - 7$$

$$= (3 \times 4) - (4) - 7$$

$$= 12 - 4 - 7$$

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

#### Solution:-

From the question it is given that m = 2

Then, substitute the value of m in the question

$$=((5 \times 2)/2)-4$$

$$=(10/2)-4$$

$$= 5 - 4$$

= 1

# 2. If p = -2, find the value of:

# (i) 4p + 7

## Solution:-

From the question it is given that p = -2

Then, substitute the value of p in the question

$$= (4 \times (-2)) + 7$$

$$= -8 + 7$$

(ii) 
$$-3p^2 + 4p + 7$$

## Solution:-

From the question it is given that p = -2

Then, substitute the value of p in the question

$$= (-3 \times (-2)^2) + (4 \times (-2)) + 7$$

$$= (-3 \times 4) + (-8) + 7$$

$$= -12 - 8 + 7$$

$$= -13$$

(iii) 
$$-2p^3 - 3p^2 + 4p + 7$$

#### Solution:-

From the question it is given that p = -2

Then, substitute the value of p in the question

$$= (-2 \times (-2)^3) - (3 \times (-2)^2) + (4 \times (-2)) + 7$$

$$= (-2 \times -8) - (3 \times 4) + (-8) + 7$$

$$= 16 - 12 - 8 + 7$$

$$= 23 - 20$$

# 3. Find the value of the following expressions, when x = -1:

## (i) 2x

**-7** 

### Solution:-

From the question it is given that x = -1

Then, substitute the value of x in the question

$$= (2 \times -1) - 7$$

$$= -2 - 7$$

+ 2

#### Solution:-

From the question it is given that x = -1

Then, substitute the value of x in the question

$$= - (-1) + 2$$

$$= 1 + 2$$

(iii) 
$$x^2 + 2x$$

## + 1 Solution:-

From the question it is given that x = -1

Then, substitute the value of x in the question

$$= (-1)^2 + (2 \times -1) + 1$$

$$= 1 - 2 + 1$$

$$= 2 - 2$$

$$= 0$$

(iv) 
$$2x^2 - x$$

## - 2 Solution:-

From the question it is given that x = -1

Then, substitute the value of x in the question

$$= (2 \times (-1)^2) - (-1) - 2$$

$$= (2 \times 1) + 1 - 2$$

$$= 2 + 1 - 2$$

4. If a = 2, b = -2, find the value of:

## Solution:-

From the question it is given that a = 2, b = -2

Then, substitute the value of a and b in the question

$$= (2)^{2} + (-2)^{2}$$
$$= 4 + 4$$
$$= 8$$

(ii) 
$$a^2 + ab$$

## + b<sup>2</sup> Solution:-

From the question it is given that a = 2, b = -2

Then, substitute the value of a and b in the question

$$= 22 + (2 \times -2) + (-2)2$$

$$= 4 + (-4) + (4)$$

$$= 4 - 4 + 4$$

$$= 4$$

## Solution:-

From the question it is given that a = 2, b = -2

Then, substitute the value of a and b in the question

$$= 2^{2} - (-2)^{2}$$

$$= 4 - (4)$$

$$= 4 - 4$$

$$= 0$$

# 5. When a = 0, b = -1, find the value of the given expressions:

### + 2b

## **Solution:-**

From the question it is given that a = 0, b = -1

Then, substitute the value of a and b in the question

$$= (2 \times 0) + (2 \times -1)$$
  
= 0 - 2  
= -2

(ii) 
$$2a^2 + b^2 + 1$$

From the question it is given that a = 0, b = -1

Then, substitute the value of a and b in the question

$$= (2 \times 0^{2}) + (-1)^{2} + 1$$
$$= 0 + 1 + 1$$
$$= 2$$

# (iii) $2a^2b + 2ab^2 +$

## ab Solution:-

From the question it is given that a = 0, b = -1

Then, substitute the value of a and b in the question

$$= (2 \times 0^2 \times -1) + (2 \times 0 \times (-1)^2) + (0 \times -1)$$

$$= 0 + 0 + 0$$

$$= 0$$

(iv) 
$$a^2 + ab$$

## + 2 Solution:-

From the question it is given that a = 0, b = -1

Then, substitute the value of a and b in the question

$$= (02) + (0 × (-1)) + 2$$
$$= 0 + 0 + 2$$
$$= 2$$

# 6. Simplify the expressions and find the value if x is equal to 2 (i) x + 7 + 4 (x - 5)

## Solution:-

From the question it is given that x = 2

We have,

$$= x + 7 + 4x - 20$$
  
=  $5x + 7 - 20$ 

Then, substitute the value of x in the equation

$$= (5 \times 2) + 7 - 20$$

$$= 10 + 7 - 20$$

$$= 17 - 20$$

$$= -3$$

(ii) 
$$3(x+2) + 5x - 7$$

From the question it is given that x = 2We have,

$$= 3x + 6 + 5x - 7$$
  
 $= 8x - 1$ 

Then, substitute the value of x in the equation

$$= (8 \times 2) - 1$$

$$= 16 - 1$$

## (iii) 6x + 5(x - 2)

#### Solution:-

From the question it is given that x = 2

We have,

$$= 6x + 5x - 10$$

$$= 11x - 10$$

Then, substitute the value of x in the equation

$$= (11 \times 2) - 10$$

$$= 22 - 10$$

## (iv) 4(2x-1) + 3x + 11

### Solution:-

From the question it is given that x = 2

We have,

$$= 8x - 4 + 3x + 11$$

$$= 11x + 7$$

Then, substitute the value of x in the equation

$$= (11 \times 2) + 7$$

$$= 22 + 7$$

## 7. Simplify these expressions and find their values if x = 3, a = -1, b =

$$-2.$$
 (i)  $3x - 5 - x + 9$ 

#### Solution:-

From the question it is given that x = 3

We have,

$$= 3x - x - 5 + 9$$

$$= 2x + 4$$

Then, substitute the value of x in the equation

$$= (2 \times 3) + 4$$

$$= 6 + 4$$

#### (ii) 2 - 8x + 4x +

### 4 Solution:-

From the question it is given that x = 3

We have,

$$= 2 + 4 - 8x + 4x$$

$$= 6 - 4x$$

Then, substitute the value of x in the equation

$$= 6 - (4 \times 3)$$

#### 1 Solution:-

From the question it is given that a = -1

We have,

$$= 3a - 8a + 5 + 1$$

$$= -5a + 6$$

Then, substitute the value of a in the equation

$$= - (5 \times (-1)) + 6$$

$$= - (-5) + 6$$

$$= 5 + 6$$

## (iv) 10 - 3b - 4 -

#### 5b Solution:-

From the question it is given that b = -2

We have,

$$= 10 - 4 - 3b - 5b$$

$$= 6 - 8b$$

Then, substitute the value of b in the equation

$$= 6 - (8 \times (-2))$$

(v) 
$$2a - 2b - 4 - 5 + a$$
  
Solution:-

From the question it is given that a = -1, b = -2We have,

$$= 2a + a - 2b - 4 - 5$$
  
 $= 3a - 2b - 9$ 

Then, substitute the value of a and b in the equation

$$= (3 \times (-1)) - (2 \times (-2)) - 9$$

$$= -3 - (-4) - 9$$

$$= -3 + 4 - 9$$

$$= -12 + 4$$

$$= -8$$

# 8. (i) If z = 10, find the value of $z^3 - 3(z - 10)$ . Solution:-

From the question it is given that z = 10 We have,

$$=z^3-3z+30$$

Then, substitute the value of z in the equation =  $(10)^3 - (3 \times 10) + 30$ 

## (ii) If p = -10, find the value of $p^2 - 2p - 100$ Solution:-

From the question it is given that p = -10 We have,

$$= p^2 - 2p - 100$$

Then, substitute the value of p in the equation

= 
$$(-10)^2 - (2 \times (-10)) - 100$$
  
=  $100 + 20 - 100$ 

# 9. What should be the value of a if the value of $2x^2 + x - a$ equals to 5, when x = 0? Solution:-

From the question it is given that x = 0We have,

$$2x^2 + x - a = 5$$
  
 $a = 2x^2 + x - 5$ 

Then, substitute the value of x in the equation

$$a = (2 \times 0^{2}) + 0 - 5$$
  
 $a = 0 + 0 - 5$   
 $a = -5$ 

## 10. Simplify the expression and find its value when a = 5 and b = -

3. 
$$2(a^2 + ab) + 3 - ab$$

Solution:-

From the question it is given that a = 5 and b = -3 We have,

$$= 2a^2 + 2ab + 3 - ab$$
  
=  $2a^2 + ab + 3$ 

Then, substitute the value of a and b in the equation

$$= (2 \times 5^{2}) + (5 \times (-3)) + 3$$

$$= (2 \times 25) + (-15) + 3$$


$$= 50 - 15 + 3$$

$$= 53 - 15$$

$$= 38$$

EXERCISE 12.4 PAGE: 246

1. Observe the patterns of digits made from line segments of equal length. You will find such segmented digits on the display of electronic watches or calculators.



If the number of digits formed is taken to be n, the number of segments required to form n digits is given by the algebraic expression appearing on the right of each pattern. How many segments are required to form 5, 10, 100 digits of the kind



#### Solution:-

(a) From the question it is given that the numbers of segments required to form n digits of the kind  $\Box$  is (5n + 1) Then,

The number of segments required to form 5 digits =  $((5 \times 5) + 1)$ = (25 + 1)= 26

The number of segments required to form 10 digits =  $((5 \times 10) + 1)$ 

$$= (50 + 1)$$

The number of segments required to form 100 digits =  $((5 \times 100) + 1)$ = (500 + 1)= 501 (b) From the question it is given that the numbers of segments required to form n digits of the kind  $\stackrel{\square}{}$  is (3n + 1)

Then,

The number of segments required to form 5 digits =  $((3 \times 5) + 1)$ = (15 + 1)

The number of segments required to form 10 digits =  $((3 \times 10) + 1)$ 

$$= (30 + 1)$$
  
 $= 31$ 

The number of segments required to form 100 digits =  $((3 \times 100) + 1)$ = (300 + 1)

(c) From the question it is given that the numbers of segments required to form n digits of the kind is (5n + 2)

The number of segments required to form 5 digits =  $((5 \times 5) + 2)$ = (25 + 2)

The number of segments required to form 10 digits =  $((5 \times 10) + 2)$ 

$$= (50 + 2)$$
  
= 52

The number of segments required to form 100 digits =  $((5 \times 100) + 1)$ 

$$= (500 + 2)$$

2. Use the given algebraic expression to complete the table of number patterns.

S.	Expression	Terms									
No.		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>		10 <sup>th</sup>		100 <sup>th</sup>	•••
(i)	2n – 1	1	3	5	7	9	-	19	-	-	-
(ii)	3n + 2	5	8	11	14	-	-	-	-	-	-
(iii)	4n + 1	5	9	13	17	-	-	-	-	-	-
(iv)	7n + 20	27	34	41	48	-	-	-	-	-	-

NCERT Solutions for Class 7 Maths Chapter 12

Algebraic Expressions

(v)	n <sup>2</sup> + 1	2	5	10	17	-	-	-	-	10001	-

**Solution:-**

(i) From the table (2n  
- 1) Then, 
$$100^{th}$$
 term =?  
Where n = 100  
=  $(2 \times 100) - 1$   
=  $200 - 1$   
= 199

(ii) From the table (3n + 2) 
$$5^{th}$$
 term =?  
Where n = 5  
=  $(3 \times 5) + 2$   
=  $15 + 2$   
=  $17$   
Then,  $10^{th}$  term =?  
Where n =  $10$   
=  $(3 \times 10) + 2$   
=  $30 + 2$   
=  $32$ 

Then, 
$$100^{th}$$
 term  
=? Where n = 100  
=  $(3 \times 100) + 2$   
=  $300 + 2$   
=  $302$ 

(iii) From the table 
$$(4n + 1) 5^{th}$$
 term =?  
Where n = 5  
=  $(4 \times 5) + 1$   
=  $20 + 1$   
=  $21$   
Then,  $10^{th}$  term =?

NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

Then, 100<sup>th</sup> term =?

Where n = 100  
= 
$$(4 \times 100) + 1$$
  
=  $400 + 1$   
=  $401$   
(iv) From the table (7n + 20) 5<sup>th</sup> term =?  
Where n = 5  
=  $(7 \times 5) + 20$   
=  $35 + 20$   
=  $55$   
Then,  $10^{th}$  term =?  
Where n =  $10$   
=  $(7 \times 10) + 20$   
=  $90$   
Then,  $100^{th}$  term =?  
Where n =  $100$   
=  $(7 \times 100) + 20$   
=  $700 + 20$   
=  $700 + 20$   
=  $700 + 20$   
=  $720$   
(v) From the table ( $n^2 + 1$ ) 5<sup>th</sup> term =?  
Where n = 5  
=  $(5^2) + 1$   
=  $25 + 1$   
=  $26$   
Then,  $10^{th}$  term =?  
Where n =  $10$   
=  $(10^2) + 1$   
=  $100 + 1$   
=  $100 + 1$ 

So the table is completed below.

## NCERT Solutions for Class 7 Maths Chapter 12 Algebraic Expressions

S.	Expression	Terms									
No.		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>		10 <sup>th</sup>		100 <sup>th</sup>	•••
(i)	2n – 1	1	3	5	7	9	-	19	-	199	-
(ii)	3n + 2	5	8	11	14	17	-	32	-	302	-
(iii)	4n + 1	5	9	13	17	21	-	41	-	401	-
(iv)	7n + 20	27	34	41	48	55	-	90	-	720	-
(v)	n <sup>2</sup> + 1	2	5	10	17	26	-	101	-	10001	-