

Case Study of the Standardized Nutrition Care Process in a Patient with Diabetes Mellitus

Andi Rezki Julianti^{1*}, Sirajuddin², Sukmawati

Correspondence email: andirezki2580@gmail.com

¹ Departement of Nutrition, Health Polytecnic of Makassar, South Sulawesi, Indonesia

ABSTRACT

In Indonesia, the number of people living with DM reached 19.5 million in 2021, making it the country with the fifth highest prevalence worldwide. This study aimed to assess the standardized nutritional care provided to a DM patient at RSAU Dr. Dody Sardjoto through an observational case study involving a single subject. Nutritional assessment indicated the patient was overweight and exhibited elevated blood pressure. Over a three-day monitoring period, nutrient intake exceeded daily requirements: energy intake ranged from 110–120%, protein 96–103%, fat 138–153%, and carbohydrates 106–128%. In conclusion, the patient was prescribed a 1500 kcal diabetic diet (DM RG). A gradual reduction in random blood glucose levels was observed, from 281 mg/dL on day one to 232 mg/dL on day three, although levels remained above the normal range

ARTICLE INFO

CASE REPORTS

Submitted: 8 July 2025

Accepted: 10 July r 2025

Keywords:

Multiple myeloma, CRAB criteria, Fracture

Copyright (c) 2024 Authors.

Access this article online

Key Messages:

- Diabetes Mellitus is a global health issue with rising prevalence, including 19.5 million cases in Indonesia as of 2021, driven by unhealthy diets and sedentary lifestyles.
- Type 2 diabetes results from insulin resistance or decreased insulin production, often linked to high sugar and fat intake and low fiber consumption, increasing the risk of complications.
- Case studies on standardized nutrition care are essential to evaluate the effectiveness of dietary interventions in managing diabetes and preventing further complications.

Introduction

Diabetes Mellitus (DM) is a degenerative disease that has long been a major health issue in both developed and developing countries. This condition is caused by various factors, one of which is lifestyle changes. DM is a metabolic disorder closely related to dietary patterns and nutritional balance. One of the main contributing factors to the development of DM is the consumption of foods high in sugar and saturated fats, along with inadequate intake of fiber and essential micronutrients.

In Indonesia, DM is also a growing public health concern. The Ministry of Health reported 19.5 million diabetes cases in 2021, representing approximately 1.5% of the total population of 172.5 million. According to WHO data, Indonesia ranks fifth globally in terms of the number of diabetes cases, following China, India, Pakistan, and the United States. In South Sulawesi alone, the number of DM cases reached 33,693 in 2018 (Risksdas, 2018).

Nutritional care for hospitalized patients with diabetes is a crucial component of intervention that must be carried out appropriately. It plays an essential role in blood glucose control and dietary management. This process includes assessment, diagnosis, intervention, monitoring, and evaluation. These stages follow a standardized nutrition care process, serving not only as a clinical protocol but also as an educational approach to foster patient commitment to dietary and lifestyle adherence. This study aims to implement the principles of the standardized nutrition care process in diabetic patients to achieve better blood glucose control.

Case Description

Assessment: Mrs. N, born November 10, 1972 aged 52 years, undergraduate education, working as a housewife, was a patient at Dr. Dody Sardjito Hospital Makassar with a medical record number 084874. Entering the hospital on March 11, 2025 in the Wren with Diabetes Milletus Media Diagnosis. Anthropometric data is 65 kg body weight, 159 cm height, body mass index 25.7 kg/m² (over body weight), ideal body weight 53.1 kg. The main complaint is weak since one day ago and lack of appetite, the history of the previous disease is hypertension and history of family diseases does not exist.

The results of laboratory tests are known to fast blood sugar is 279 mg/dl (height) compared to normal references 70-126 mg/dl. Blood sugar 2 hours PP is 419 mg/dl (tall) compared to the reference 200 mg/dl. Triglyceride 235 mg/dl (high) compared to the reference (200 mg/dl). Blood pressure examination on March 12, 2025 is 160/100 mmHg (hypertension) compared to normal references 120/80 mmHg. Body temperature 36 C (normal), pulse 90 times/minute (normal) compared to the reference 60-100 times/minute. Respiration rate is 20 times/minute (normal) compared to the reference (16-20 times per minute).

The patient's dietary history before entering the hospital is eating at home 2 times a day and 1 time interrupting with staple white rice, animal side dishes of fish and cooking chickens or fried, vegetable side dishes of orek tempeh and sometimes fried flour, often consume vegetables and fruits such as coconuts. Often consume fried food and fruit ice. Recall consumption results are known to be 1177.4 kcal while their needs are 1525, 9 kcal or 77% reach (less) needs. Protein intake is 49.5 g while the need is 76.2 g or 64% achieves (less) needs, fat intake is 13.8 g, 33.9 g or 40% of the achievement of lack of need, carbohydrate intake 216.6 g, needs 228.8 g or 94% (sufficient).

Nutritional diagnosis; NI-2.1 Lack of oral food and drink intake is related to lack of appetite characterized by food intake below normal, namely energy (70%), protein (58%), fat (36%). NI-5.4 Decreased need for sodium nutrients associated with increased blood pressure is characterized by blood pressure 160/100 mmHg. NC-2.2 Changes in Laboratory Value Related to Nutrition associated with metabolic disorders are characterized by the results of the examination indicating GDP levels 279 mg/dl, GD2JPP 419 mg/dl and triglycerides 235 mg/dl.

Discussion

Nutritional intervention. Nutritional needs are determined based on the consensus of Perkeni, 2021, namely ideal body weight multiplied by 25 kcal with age correction (BMR-5%), activity correction (age correction + 10%), metabolic stress correction (activity correction + 10%). The total energy requirement is 1525.9 kcal. Protein needs are set 20% of energy requirements or 76.2 g. Fat requirements are set at 20% of energy requirements or 33.9 g. Carbohydrate requirements are 60% of energy requirements or 228.8 g.

Nutrition therapy given to patients is a type of DM 1500 low salt diet 1. Soft food form, oral administration method, 3 times the main food and 1 time interlude food. Oral giving route. The purpose of nutritional therapy is to meet the patient's nutritional needs, achieve and maintain normal sugar levels, reach normal blood pressure.

Diet Requirements: Energy As needed from the calculation results, protein 76.2 gr/day. Fat 33.9 gr/day, carbohydrates 228.8 gr/day, sodium intake is limited to 600 mg/day, the use of pure sugar in drinks and food is not allowed unless the sugar is slightly as a spice. Fiber intake is recommended 25 gr/hr by prioritizing water soluble fiber contained in vegetables and fruit. Enough vitamins and minerals. Feeding pays attention to 3 J (amount, type, schedule), limiting consumption of saturated fat and cholesterol. This diet requirement has been met after intervention for three days.

Monitoring and evaluation. Food assessment intake is carried out for 3 days, namely on March 13-15, 2025 in the inpatient room (Wren) RSAU Dr. Dody Sardjoto. The food served to patients is hospital food. The food served is adjusted to the calculation of the patient's nutritional needs. Eating intake that is observed 3 days includes breakfast, lunch, and dinner.

Table 1. Laboratory examination results

Date Intervention	Energy and Macronutrient	RDA	Actual intake	Percentage of RDA	Status
13 March 2025	Energy (kcal)	1525,9	1685,9	110%	Optimal
	Protein (g)	76,2	77,7	101%	Optimal
	Fat (g)	33,9	47	138%	Over
	Carbohydrate (g)	228,8	242,7	106%	optimal
14 March 2025	Energy (kcal)	1525,9	1699,4	111%	optimal
	Protein (g)	76,2	74	96%	Optimal
	Fat (g)	33,9	47,6	140%	Over
	Carbohydrate (g)	228,8	248,7	108%	Optimal
15 March 2025	Energy (kcal)	1525,9	1835,7	120%	Optimal
	Protein (g)	76,2	79,1	103%	Optimal
	Fat (g)	33,9	51,9	153%	Over
	Carbohydrate (g)	228,8	295	128%	over

The following is the image of the radiology examination results.




Figure 1. Cranium AP Lateral (Conventional non- contrast)

Conclusion

Based on the case description, the CRAB criteria approach in diagnosing multiple myeloma indicated that this patient meets the criteria for multiple myeloma. The CRAB criteria can be used as a guideline for diagnosing multiple myeloma, especially in areas with limited access to comprehensive laboratory testing. This approach enables a quick diagnosis, allowing prompt management of the condition.

Funding: This research received no external funding

Acknowledgments: The authors would like to thank RSUD Provinsi Banten for providing access to medical records to support this case report

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Sutandyo N, Firna E, Agustina J, Prayogo N, Widjaja L. Clinicopathology Profile and Bone Involvement of Multiple Myeloma Patients in Dharmas National Cancer Hospital, Indonesia. *Asian Pacific Journal of Cancer Prevention*. 2015 Oct 6;16(15):6261–5.
2. An ZY, Fu H, He Y, Zhu X, Huang QS, Wu J, et al. Projected Global Trends in Hematological Malignancies: Incidence, Mortality, and Disability-Adjusted Life Years from 2020 to 2030. *Blood*. 2023 Nov 2;142(Supplement 1):3810–3810. Cowan AJ, Allen C, Barac A, Basaleem H, Bensenor I, Curado MP, et al. Global Burden of Multiple Myeloma. *JAMA Oncol*. 2018 Sep 1;4(9):1221.
3. Fazio M, Del Fabro V, Parrinello NL, Allegra A, Markovic U, Botta C, et al. Multiple Myeloma in 2023 Ways: From Trials to Real Life. *Current Oncology*. 2023 Nov 3;30(11):9710–33.
4. Cowan AJ, Allen C, Barac A, Basaleem H, Bensenor I, Curado MP, et al. Global Burden of Multiple Myeloma. *JAMA Oncol*. 2018 Sep 1;4(9):1221.
5. Palumbo A, Avet-Loiseau H, Oliva S, Lokhorst HM, Goldschmidt H, Rosinol L, et al. Revised International Staging System for Multiple Myeloma: A Report From International Myeloma Working Group. *Journal of Clinical Oncology*. 2015 Sep 10;33(26):2863–9.
6. Palumbo A, Anderson K, Battista SG. Medical Progress Multiple Myeloma.

7. Rajkumar SV. Multiple myeloma: 2022 update on diagnosis, risk stratification, and management. *Am J Hematol*. 2022 Aug 23;97(8):1086–107.
8. Kumar SK, Rajkumar V, Kyle RA, et al. Multiple myeloma. *Nat Rev Dis Primers*. 2017;3:17046. DOI: 10.1038/nrdp.2017.46
9. Gau YC, Yeh TJ, Hsu CM, Hsiao SY, Hsiao HH. Pathogenesis and Treatment of Myeloma-Related Bone Disease. *Int J Mol Sci*. 2022;23(6):3112. Published 2022 Mar 14. doi:10.3390/ijms23063112
10. Teramachi J, Miki H, Nakamura S, Hiasa M, Harada T, Abe M. Myeloma bone disease: pathogenesis and management in the era of new anti-myeloma agents. *J Bone Miner Metab*. 2023;41(3):388-403. doi:10.1007/s00774-023-01403-4
11. Facon T, Kumar S, Plesner T, et al. Daratumumab plus Lenalidomide and Dexamethasone for Untreated Myeloma. *N Engl J Med*. 2019;380(22):2104-2115. DOI: 10.1056/NEJMoa1817249
12. Annadatha A, Acharya S, Shukla S, Bhawane A, Varma A, Andhale A. Multiple Myeloma presenting as Hypercalcemic Crisis: A case report. *Medical Science*. 2020;24(105):3148-3152
13. Costa LJ, Chhabra S, Godby KN, et al. Daratumumab, Carfilzomib, Lenalidomide and Dexamethasone (Dara-KRd) Induction, Autologous Transplantation and Post-Transplant, Response-Adapted, Measurable Residual Disease (MRD)-Based Dara-Krd Consolidation in Patients with Newly Diagnosed Multiple Myeloma (NDMM). *Blood*. 2019;134(Supplement_1):860. DOI: 10.1182/blood-2019-123170
14. Raje N, Berdeja J, Lin Y, et al. Anti-BCMA CAR T-Cell Therapy bb2121 in Relapsed or Refractory Multiple Myeloma. *N Engl J Med*. 2019;380(18):1726-1737. DOI: 10.1056/NEJMoa1817226