



February 8, 2023

Golden Horseshoe Cycling Hub Concussion Policy

Preamble

1. This Policy is based on the 5th Consensus Statement on Concussion in Sport that was released in April 2017. This Policy interprets the information contained in the report that was prepared by the 2017 Concussion in Sport Group (CISG), a group of sport concussion medical practitioners and experts, and adapts concussion assessment and management tools.
2. This Policy is intended to be compliant with Ontario's Rowan's Law (Concussion Safety), 2018. If any provision of the policy is in conflict with Rowan's Law, the legislation shall take precedence.
3. The CISG suggested 11 'R's of Sport-Related Concussion ("SRC") management to provide a logical flow of concussion management. This Policy is similarly arranged. The 11 R's in this Policy are: Recognize, Remove, Re-Evaluate, Rest, Rehabilitation, Refer, Recover, Return to Sport, Reconsider, Residual Effects, and Risk Reduction.
4. A concussion is a clinical diagnosis that can only be made by a physician.

Definitions

5. The following terms have these meanings in this Policy:
 - a. "Organization" – Golden Horseshoe Cycling Hub
 - b. "Participant" – Coaches, athletes, volunteers, officials and other Registered Individuals
 - c. "Registered Individuals" – All individuals employed by, or engaged in activities with the Organization, including but not limited to, employees, volunteers, administrators, committee members and directors and officers.
 - d. "Suspected Concussion" – means the recognition that an individual appears to have either experienced an injury or impact that may result in a concussion or who is exhibiting unusual behaviour that may be the result of concussion.
 - e. "Sport-Related Concussion ("SRC") – A sport-related concussion is a traumatic brain injury induced by biomechanical forces. Several common features that may be used to define the nature of a SRC may include:
 - i. Caused either by a direct blow to the head, face, neck or elsewhere on the body with an impulsive force transmitted to the head.
 - ii. Typically results in the rapid onset of short-lived impairment of neurological function that resolves spontaneously. However, in some cases, signs and symptoms evolve over a number of minutes to hours.
 - iii. May result in neuropathological changes, but the acute clinical signs and symptoms largely reflect a functional disturbance rather than a structural injury and, as such, no abnormality may be visibly apparent.
 - iv. Results in a range of clinical signs and symptoms that may or may not involve loss of consciousness. Resolution of the clinical and cognitive



February 8, 2023

features typically follows a sequential course. However, in some cases symptoms may be prolonged.

Purpose

6. The Organization is committed to ensuring the safety of Participants in its activities. The Organization recognizes the increased awareness of concussions and their long-term effects and believes that prevention of concussions is paramount to protecting the health and safety of Participants.
7. This Policy describes the common signs and symptoms of a concussion and how to identify them, the protocol to be followed in the event of a possible concussion, and a Return to Sport protocol should a concussion be diagnosed. Awareness of the signs and symptoms of concussion and knowledge of how to properly manage a concussion is critical to recovery and helping to ensure the individual is not returning to physical activities too soon, risking further complication.
8. This Policy applies to all activities and events for which the Organization is the governing or sanctioning body including, but not limited to, competitions, practices, and training sessions.

Registration

9. When an individual under the age of 26 years old registers with the Organization, the individual must provide written or electronic confirmation that they have reviewed concussion awareness resources within the past 12 months. The Ontario Government has produced age-appropriate concussion resources located here:
 - a. Ages 10 and under
 - b. Ages 11-14
 - c. Ages 15+
10. Individuals under the age of 26 years old must also sign the Concussion Code of Conduct (Appendix A).
11. For athletes younger than 18 years old, the athlete's parent or guardian must also provide confirmation that they have also reviewed the concussion resources as well and signed the Concussion Code of Conduct.
12. Coaches, officials and team trainers must provide confirmation that they have also reviewed the concussion resources and sign the Concussion Code of Conduct; but not if they will be interacting exclusively with athletes who are 26 years old or older.

Recognizing Concussions

13. If a Participant demonstrates or reports any of the following red flags, an on-site licensed healthcare professional shall be summoned and, if deemed necessary, an ambulance should be called¹:
 - a. Neck pain or tenderness

¹ If an onsite healthcare professional is not available, an ambulance should be called.





February 8, 2023

- b. Double vision
 - c. Weakness or tingling / burning in arms or legs
 - d. Severe or increasing headache
 - e. Seizure or convulsion
 - f. Loss of consciousness
 - g. Deteriorating conscious state
 - h. Vomiting more than once
 - i. Increasingly restless, agitated, or combative
 - j. Increased confusion
14. The following observable signs may indicate a possible concussion:
- a. Lying motionless on the playing surface
 - b. Slow to get up after a direct or indirect hit to the head
 - c. Disorientation or confusion / inability to respond appropriately to questions
 - d. Blank or vacant look
 - e. Balance or gait difficulties, absence of regular motor coordination, stumbling, slow laboured movements
 - f. Facial injury after head trauma
15. A concussion may result in the following symptoms:
- a. Headache or "pressure in head"
 - b. Balance problems or dizziness
 - c. Nausea or vomiting
 - d. Drowsiness, fatigue, or low energy
 - e. Blurred vision
 - f. Sensitivity to light or noise
 - g. More emotional or irritable
 - h. "Don't feel right"
 - i. Sadness, nervousness, or anxiousness
 - j. Neck pain
 - k. Difficulty remembering or concentrating Feeling slowed down or "in a fog"
16. Failure to correctly answer any of these memory questions may suggest a concussion:
- a. What venue are we at today?
 - b. Where was your last major competition?
 - c. What day is it?
 - d. What event are you participating in?

Removal from Sport Protocol

17. In the event of a Suspected Concussion where there are observable signs of a concussion, symptoms of a concussion, or a failure to correctly answer memory questions, the Participant should be immediately removed from participation by a designated person who is either an on-site Organization staff member and/or Coach, Volunteer, or Contracted First Aid or Medical company.
18. After removal from participation, the following actions should be taken:



February 8, 2023

- a. The designated person who removed the Participant should consider calling 9-1-1;
 - b. The Organization must make and keep a record of the removal;
 - c. The designated person must inform the Participant's parent or guardian if the Participant is younger than 18 years old, and the designated person must inform the parent or guardian that the Participant is required to undergo a medical assessment by a physician or nurse practitioner before the Participant will be permitted to return to participation; and
 - d. The designated person will remind the Participant, and the Participant's parent or guardian as applicable, of the Organization's Return-to-Sport protocol as described in this Policy.
19. Participants who have a Suspected Concussion and who are removed from participation should:
- a. Be isolated in a dark room or area and stimulus should be reduced
 - b. Be monitored
 - c. Have any cognitive, emotional, or physical changes documented
 - d. Not be left alone (at least for the first 1-2 hours)
 - e. Not drink alcohol
 - f. Not use recreational/prescription drugs
 - g. Not be sent home by themselves
 - h. Not drive a motor vehicle until cleared to do so by a medical professional
20. A Participant who has been removed from participation due to a suspected concussion should not return to participation until the Participant has been assessed medically by a physician or nurse practitioner who is familiar with the Sport Concussion Assessment Tool – 5th Edition (SCAT5) (for Participants over the age of 12) or the Child SCAT5 (for Participants between 5 and 12 years old), even if the symptoms of the concussion resolve.

Re-Evaluate

21. A Participant with a Suspected Concussion must be evaluated by a licensed physician or nurse practitioner who should conduct a comprehensive neurological assessment of the Participant and determine the Participant's clinical status and the potential need for neuroimaging scans.

Rest and Rehabilitation

22. Participants with a diagnosed SRC should rest during the acute phase (24-48 hours) but can gradually and progressively become more active so long as activity does not worsen the Participant's symptoms. Participants should avoid vigorous exertion.
23. Participants must consider the diverse symptoms and problems that are associated with SRCs. Rehabilitation programs that involve controlled parameters below the threshold of peak performance should be considered.

February 8, 2023

Refer

24. Participants who display persistent post-concussion symptoms (i.e., symptoms beyond the expected timeline for recovery – 10-14 days for adults and 4 weeks for children) should be referred to physicians with experience handling SRCs.

Recovery and Return to Sport

25. SRCs have large adverse effects on cognitive functioning and balance during the first 24-72 hours after injury. For most Participants, these cognitive defects, balance and symptoms improve rapidly during the first two weeks after injury. An important predictor of slower recovery from an SRC is the severity of the Participant's initial symptoms following the first few days after the injury.
26. The table below represents a graduated return to sport for most Participants, in particular those that did not experience high severity of initial symptoms after the following the first few days after the injury.

Stage	Aim	Activity	Stage Goal
1	Symptom-limited activity	Daily activities that do not provoke symptoms	Gradual reintroduction of activities
2	Light aerobic exercise	Walking or stationary cycling at slow to medium pace. No resistance training	Increase heart rate
3	Sport-specific exercise	Light drills. No head impact activities	Add movement
4	Non-contact training drills	Harder training drills. May start progressive resistance training	Exercise, coordination and increased thinking
5	Full contact practice	Following medical clearance, participate in normal training activities	Restore confidence and assess functional skills by coaching staff
6	Return to sport	Normal participation	

Table 1 – Return to Sport Strategy

27. An initial period of 24-48 hours of both physical rest and cognitive rest is recommended before beginning the Return to Sport strategy.
28. There should be at least 24 hours (or longer) for each step. If symptoms reoccur or worsen, the Participant should go back to the previous step.
29. Resistance training should only be added in the later stages (Stage 3 or Stage 4).
30. If symptoms persist, the Participant should return to see a physician.

February 8, 2023

31. The Participant's Return-to-Sport strategy should be guided and approved by a physician with regular consultations throughout the process.
32. The Participant must provide the Organization with a medical clearance form, signed by a physician, following Stage 5 and before proceeding to Stage 6.

Reconsider

33. The 2017 Concussion in Sport Group (CISG) considered whether certain populations (children, adolescents, and elite athletes) should have SRCs managed differently.
34. It was determined that all Participants, regardless of competition level, should be managed using the same SRC management principles.
35. Adolescents (13 to 18 years old) and children (5 to 12 years old) should be managed differently. SRC symptoms in children persist for up to four weeks. More research was recommended for how these groups should be managed differently, but the CISG recommended that children and adolescents should first follow a Return to School strategy before they take part in a Return to Sport strategy. A Return to School strategy is described below.

Stage	Aim	Activity	Stage Goal
1	Daily activities at home that do not give the child symptoms	Typical activities of the child during the day as long as they do not increase symptoms (e.g., reading, texting, screen time). Start with 5–15 min at a time and gradually build up	Gradual return to typical activities
2	School activities	Homework, reading or other cognitive activities outside of the classroom	Increase tolerance to cognitive work
3	Return to school part-time	Gradual introduction of schoolwork. May need to start with a partial school day or with increased breaks during the day	Increase academic activities
4	Return to school full time	Gradually progress school activities until a full day can be tolerated	Return to full academic activities and catch up on missed work

Table 2 – Return to School Strategy



February 8, 2023

Residual Effects

36. Participants should be alert for potential long-term problems such as cognitive impairment and depression. The potential for developing chronic traumatic encephalopathy (CTE) should also be a consideration, although the CISG stated that “a cause-and-effect relationship has not yet been demonstrated between CTE and SRCs or exposure to contact sports. As such, the notion that repeated concussion or subconcussive impacts cause CTE remains unknown.”

Risk Reduction and Prevention

37. The Organization recognizes that knowing a Participant's SRC history can aid in the development of concussion management and the Return to Sport strategy. The clinical history should also include information about all previous head, face, or cervical spine injuries. The Organization encourages Participants to make coaches and other stakeholders aware of their individual histories.

Non-Compliance

38. Failure to abide by any of the guidelines and/or protocols contained within this policy may result in disciplinary action in accordance with the Organization's policies for discipline and complaints.

Liability

39. The Organization shall not be liable for any Participant or other individual's use or interpretation of this Policy. Further, none of the Organization's members, directors, officers, employees, agents, representatives and other individuals involved in any way in the administration of this Policy shall be liable to any other individual in any way, in relation to any lawful acts or omissions committed in the honest application, administration, and/or enforcement of this Policy.