

BCFRAG expert team notes

Notes for meeting 15 May 2024 <https://indico.cern.ch/event/1412563>

Minutes online –

- Participants:
 - Andreas Meyer
 - Adrián Irles
 - Eli Ben Haim
 - Paolo Azzurri
 - Andrzej Siodmok
 - Loukas Gouskos
 - Torbjörn Sjöstrand
 - Keisho Hidaka
 - German Rodrigo
 - Maria Ubiali
- Synergies with W-mass (FT document 4.1) color reconnection (Raimund Stroemer,).
 - Clarify whether we can use LEP data with newer algorithms to see possible effects in the data.
 - Try also with simulations of new data
- Torbjorn:
 - ECFA ee study workshop, comparison of color reconnection models, a lot of development also in pp, but not so many news in pythia.
 - Previous reports have been extensive on the theory side. Maybe good to be more experimental.
- Andrzej: HERWIG studies, q/g separation in Les Houches (mostly for pp collisions)
- Loukas:
 - also for flavour tagging, and g/q separation, test different models and parameters, also for hadronization, also for different jet identification algorithms => make a breakdown of variations (model and exp. parameters) and their impact on key measurements such as $H \rightarrow qq$ - initially on simulations
 - In the Les Houches context, there is plans to have results in the coming months.
 - CERN fellow working on LEP data now, converted into EDM4hep format
 - Overlap with CEPC efforts, Manqi et al ? contact Manqi again (also ask about Z coupling)
 - Andrzej : <https://inspirehep.net/literature/1591528>
- Andrzej: G.Salam, A.Mitov are working on jet flavour algorithms.
<https://arxiv.org/abs/2205.11879> <https://arxiv.org/abs/2208.11138>
<https://arxiv.org/abs/2306.07314>

- Detector performance: Loukas, studies ongoing, how much would observables be affected when degrading detectors. Studies of simulation and detector groups $\Leftrightarrow$ cross coordinate.
- Andrzej: ML approaches exist, but are still in the beginning.
- Torbjorn: naive assumption that what happens inside jet cone factorizes from the rest of the event is not true.
- Andrzej and Torbjorn: analytical calculations: b quark production and fragmentation are independent, No tracing of origin, only momentum. As a result, fragmentation function (must be) independent. In contrast in HERWIG or PYTHIA, depending how the neighbourhood looks, end up with a different hadron spectrum.
- Keisho: Migration matrix analysis (by Manqi for CEPC) should be compared between CEPC and FCC. Loukas: such analysis exists at Z pole. Single W production in ee around 140 GeV has been studied.

Previous meetings:

<https://indico.cern.ch/event/1332815/>

<https://indico.cern.ch/event/1318673/>

Previous notes (copied from here):

<https://docs.google.com/document/d/1WqW--pewiVoS0UNkj0Pqe-3x2bdq9PIbQevTKszsG4w/edit>

BCFrag: Measurement of b- and c-fragmentation functions and hadronisation rates

What?

- Insufficient knowledge of hadronisation and fragmentation of heavy quarks could introduce significant systematic uncertainties $\rightarrow$ for example in AFB (b/c)
arXiv:2010.08604

Theory and pheno targets

- Are existing hadronization models (strong fragmentation, cluster fragmentation) flexible enough, or do we need new ideas?
- Identify calibration observables that are well understood theoretically and unaffected by BSM physics.
- See also
https://indico.desy.de/event/33640/contributions/130328/attachments/77658/100472/ECFA22_Hadronization.pdf

Theory and pheno targets:

- Event shapes, angular distributions
- Fragmentation functions
- Specific observables, i.e.
 - mW,

- AFB(b,c) (at Z-pole and above)

Gsplit Measurement of gluon splitting to bb / cc, interplay with separating $h \rightarrow$ gluons from $h \rightarrow$ bb/cc

What?

- Insufficient knowledge of gluon splitting to heavy quarks could introduce significant systematic uncertainties in precision Higgs and electroweak measurement

Theory and pheno targets

- How to consistently implement gluon splitting in parton shower tools (modelling and free parameters)

target physics observables

- 3/4-jet events with multiple heavy quark ids.
- Jet substructure observables with hadron tags inside jet.

#####

Minutes of first meeting, 2023/08/31

Participants:

Eli Ben-Haim <benhaim@in2p3.fr>

Maria Ubiali <mu227@cam.ac.uk>

Andrzej Siodmok <a.siodmok@cern.ch>

simon.plaetzer@uni-graz.at <simon.plaetzer@uni-graz.at>

Loukas Gouskos <Loukas.Gouskos@cern.ch>

Torbjörn Sjöstrand <torbjorn.sjostrand@hep.lu.se>

&

Adrián Irlés

Andreas Meyer

Ayres Freitas

Paolo Azurri

Fabio Maltoni

Notes:

Torbjorn - discussing slides 6/7 of his presentation: Fragmentation cannot be described as a convolution of a perturbative matrix element and a fragmentation function $f(z)$ with $0 < z < 1$. In string fragmentation, color strings stretching between partons cause acceleration and produce a *hardening* of the hadron spectra at high energies.

Needs fragmentation model. There will still be free parameters, which we may have to determine ad hoc.

Ayres - Q: Are the currently used frameworks flexible enough (after adding more mesons than currently implemented in e.g. Pythia) to be precise/accurate enough for observables with sub-permille precision?

Simon: maybe rather ask can we reach accuracy ?

Ayres - Q: Are fragmentation functions universal, or do they depend on types of observables?

Eli: but parameters will not be universal.

Simon: modeling developments in disentangling hadronisation and fragmentation.
Cross-talk between parton shower and fragmentation. New work needed for NLL accurate showers.

Paolo - Q: Can the LHC data be used to test new and more accurate models in a way that help for the future e+e- measurements ?

Torbjorn: The environment is quite different. Charm and bottom production rates at LHC are not precisely known.

do we know the total c and production rate?

can we trust the extracted fragmentation functions ?

But something could maybe still be done (e.g. amount of non-B hadron activity in a cone around a B hadron).

Ayres -Q: is gluon splitting and quark fragmentation fundamentally different, conceptually ?

- Sjostrand:

Gluon splitting into cc/bb only in the perturbative (parton shower) region, not in the string/cluster fragmentation. But need to ask how to account for charm/bottom masses in the shower.

- in the string model, the nonpert. step does not add additional c or b.
everything is from pert. stage.

- one example of an uncertainty is alphas, where the in the limit of $p_t=0$, heavy quarks can still be treated, but for gluons everything is different.

Simon: Do we need to consider multi-parton fragmentation functions? It could be formulated in a theoretically rigorous framework, with factorization, but may be very complicated.

Useful data input: correlations among hadrons, Lund-plane observables

Transition shower to hadronization is a fuzzy one.

- we are at the start of exploring fragmentation/hadronisation, effect
- use LHC data, e.g. ALICE to test correlations between different heavy quarks.

(small-angle

Adrian:

- make a list of relevant observables (for the future ee collider) for which these things are relevant.

#####

Examples of observables (list made by Torbjörn Sjöstrand)

Examples of Bfrag/Gsplit observables:

Approximately ordered in terms of increasing differentiation, as viewed by a theorist, disregarding experimental complications. In reality, several measurements would be intertwined.

"EpT" will be used as shorthand for E or pT hardness variable of particle, the former normally for e+e- and the latter for pp.

* Inclusive B/D particle (mesons + baryons) production cross section

- e+e-: primary production well known from theory, so "excess" is from gluon splitting
- pp: combines primary production, gluon splitting, and MPI (multiparton interactions) contributions, each with significant theoretical uncertainties

* Flavour composition, as far back in decay chains as can be traced

(even equal D^0 and D^{*+} rates gives unequal D^0 and D^+ ones)

- e+e-: we do not expect sizeable momentum dependence, but interesting to contrast mesons and baryons for smaller ones
- pp: significant pT dependence observed and to be studied further, also high- vs. low-multiplicity events, rapidity, ..., which is important for development/tuning of colour reconnection models

* Particle-antiparticle production asymmetries

- e+e-: none expected, except tiny from CP-violation in oscillations
- pp: asymmetries expected and observed from p flavour content, increasing at larger rapidities; relates to how string (and cluster?) fragmentation connects central rapidities to beam remnants

* Momentum spectra

- e+e-: dn/dx_E where $x_E = 2E_{\text{had}} / E_{\text{cm}}$; basic distribution for tuning of "fragmentation function"
- pp: dn/dp_T and dn/dy give basic production kinematics, but the many production channels gives less easy interpretation

* Energy flow around B/D hadron, excluding this hadron itself, as a test that dead cone effects are correctly described

- e+e-: $dE/d\theta$ where θ is distance from B/D on the sphere
- pp: dp_T/dR where R is distance in (η , ϕ) or (y , ϕ) space, only applied for B/D above some pT threshold

* B/D hadron fraction of total E or pT in a jet, with

$x = E_{pT_had} / E_{pT_jet}$, as a test of the fragmentation function combined with almost collinear radiation, suitably for some slices of EpT

- e+e-: draw a jet cone in theta around B/D and measure x
- pp: draw a jet cone in R around B/D and measure x

* **As above, but with a veto that no other B/D** should be inside the jet cone, so as to suppress the gluon splitting contribution

* Distribution in number of reconstructed B/D hadrons, as a measure of how often several pairs are produced

* Separation inside B/D pairs, where large separation suggests back-to-back primary production, while small separation suggests gluon splitting

- e+e-: separation in theta
- pp: separation both in phi and in R, since for primary production $\phi = \pi$ is hallmark with eta/y separation less interesting, while gluon splitting means R is small while phi and y/eta individually are less interesting

* Hardness difference within (reasonably hard) pairs,
 $\Delta = (E_{pT_max} - E_{pT_min}) / (E_{pT_max} + E_{pT_min})$,
 where for gluon splitting $x^2 + (1 - x)^2$ translates to $1 + \Delta^2$

- e+e-: separately for small or large theta
- pp: separately for large or small phi

* For a pair with small separation, say $\theta/R < 0.7$, draw a cone around the midpoint of the two, say again $\theta/R = 0.7$, and find the fraction $x = (E_{pT_had1} + E_{pT_had2}) / E_{pT_jet}$, to quantify loss to showers and hadronization. This loss would be reduced if colour reconnection often would make the b-bbar or c-cbar into a singlet, rather than the default octet where the two fragment separately.

* In events with two B/D pairs, many observables become possible. There are four possible particle-antiparticle pairs (more if B-Bbar mixing is taken into account), each of which can be studied according to the two points above. In addition, a pair with a small separation would suggest a gluon splitting, while one with a large ditto a primary production. For pp, two back-to-back pairs would suggest MPI. One can try to classify events into most likely history and study relative composition of

- two separate hard processes (MPIs, pp only),
- one hard process and one gluon split,
- two gluon splits on same side of the event,
- two gluon splits on opposite sides.

* Even if one B/D is missed in pp, so that only three B/D are observed, one can study the three pairings, and see whether either pair has a small R or a large phi. Again relative rates will

provide info on the composition of production mechanisms.

#####

Minutes of second meeting, 2023/10/03

Private overleaf document to be filled

→ Link in the agenda of the event. [Please read it and add suggestions/corrections wherever needed.](#)

First draft of the document started. Current approach:

- identify the main analysis/physics cases that will suffer the most from uncertainties on BCfrag/GSplit.
- Describe the existing tools (experimental and theory)
- Find a set of observables that could be used as entry point for
 - experimentalists to study new detector models/reconstruction tools
 - Theorists to test new models
 - Use LEP data to validate these new models?

Discussion

- **Ubiali/Platzer discussion** (sorry, I missed most of it due to unexpected construction works for ~5minutes...)
- Definition of observables
 - Should start with a short list for the ECFA-Paestum workshop timescale in mind.
 - Further in time, ideally, we find people who are interested in reproducing these observables and study them for different models/tunes.
 - Theory and/or full simulation studies? LEP data?
- Using LEP data:
 - Simon has some experience: it is very useful but very technically challenging. It requires an expert from LEP to understand the data and how to use it.
 - Loukas CERN team will hire a technical student to look into that together with ALEPH physicists. The idea would be to migrate these data to the Key4HEP (or similar) format.
 - Comment: it could be interesting to do the same for other experiments.
- ECFA-Paestum meeting
 - We will share the zoom link.
 - Overlap with the TwoF focus topic.. Which is highly correlated with this one. We will contact the conveners and try not to make them overlap.