

References

Pistone et al 2017 Molecular phylogeny of bark and ambrosia beetles (Curculionidae: Scolytinae) based on 18 molecular markers

Hulcr et al. 2010 "Repeated evolution of crop theft in fungus-farming ambrosia beetles." *Evolution* 64.11 (2010): 3205-3212. Maksoud, S., Granier, B.R. and Azar, D., 2022. Palaeoentomological (fossil insects) outcrops in Lebanon. *Carnets Geol.*, 22(16), pp.699-743.

El Hajj, Layla, et al. "Dysodiles from the lower Barremian of Lebanon: Insights on the fossil assemblages and the depositional environment reconstruction." *Cretaceous research* 120 (2021): 104732.

Diel 2022 First experimental evidence for active farming in ambrosia beetles and strong heredity of garden microbiomes <https://royalsocietypublishing.org/doi/10.1098/rspb.2022.1458>

Scott 2008 Bacterial Protection of Beetle-Fungus Mutualism

Trapp, Susan, and Rodney Croteau. "Defensive resin biosynthesis in conifers." *Annual review of plant biology* 52.1 (2001): 689-724

Azar, Dany. "Lebanese amber: a "Guinness Book of Records". " *Annales Universitatis Paedagogicae Cracoviensis. Studia ad Didacticam Biologiae Pertinentia* 2.I (2012): 44-60

Azar, Dany. "Preservation and accumulation of biological inclusions in Lebanese amber and their significance." *Comptes Rendus Palevol* 6.1-2 (2007): 151-156

Boderau et al 2024 The first aquatic insect fossil from the Cenomanian Konservat-Lagerstätte of Haql (Lebanon) fills a gap in the fossil record of giant water bugs (Heteroptera: Belostomatidae). *Zootaxa* 5562: 1. 11-22 Dec.

NPR 2024 In Lebanon, the 'Amber Man' digs up golden time capsules from the age of the dinosaurs <https://www.npr.org/2024/06/28/g-s1-5121/lebanon-ancient-amber-cretaceous-dinosaurs-flowers>

Peris, D et al 2021 Origin and evolution of fungus farming in wood-boring Coleoptera—a palaeontological perspective. *Biological Reviews*, 96(6), 2476-2488.

Seyfullah et al 2020 Revealing the diversity of amber sourceplants from the Early Cretaceous Crato Formation, Brazil

Langenheim, J. H. (1995). Biology of amber-producing trees: focus on case studies of *Hymenaea* and *Agathis*.

Mayers et al ,2020.Patterns of coevolution between ambrosia beetle mycangia and the Ceratocystidaceae, with five new fungal genera and seven new species. *Persoonia-Molecular Phylogeny and Evolution of Fungi*, 44(1), 41-66.

Peris, D., 2020. Coleoptera in amber from Cretaceous resiniferous forests. *Cretaceous Research*, 113, p.104484.

Delclòs, 2023 "Amber and the Cretaceous resinous interval." *Earth-Science Reviews* (2023): 104486.

Cockx, P., & McKellar, R. C. (2024). Bonebed amber deposits: a review of taphonomy and palaeontological significance. *Evolving Earth*, 100032.

McCoy, V. E., Barthel, H. J., Boom, A., Peñalver, E., Delclòs, X., & Solórzano-Kraemer, M. M. (2021). Volatile and semi-volatile composition of Cretaceous amber. *Cretaceous Research*, 127, 104958

Beutel, R.G., Xu, C., Jarzembowski, E., Kundrata, R., Boudinot, B.E., McKenna, D.D. and Goczał, J., 2024. The evolutionary history of Coleoptera (Insecta) in the late Palaeozoic and the Mesozoic. *Systematic Entomology*.

ńczak, J., Kosakowski, P., Drzewicz, P. and Zakrzewski, A., 2024. Fossil resins—A chemotaxonomical overview. *Earth-Science Reviews*, p.104734.

Poinar, G. O., & Mastalerz, M. (2000). Taphonomy of fossilized resins: determining the biostratigraphy of amber. *Acta Geologica Hispanica*, 171-182.

Ulyshen, Michael D. "Wood decomposition as influenced by invertebrates." *Biological Reviews* 91.1 (2016): 70-85.

Chen, X. (2019). A new genus and species of Lymexylidae (Insecta: Coleoptera) from mid-Cretaceous amber of northern Myanmar. *Paleontological Journal*, 53, 1203-1205.

Wolf-Schwenninger, K. (2011). The oldest fossil record of Lymexylidae (Insecta: Coleoptera) from the Lower Cretaceous Crato formation of Brazil. *Insect Systematics & Evolution*, 42(2), 205-212.

Kirejtshuk, A. G., Azar, D., Beaver, R. A., Mandelshtam, M. Y., & Nel, A. (2009). The most ancient bark beetle known: a new tribe, genus and species from Lebanese amber (Coleoptera, Curculionidae, Scolytinae). *Systematic Entomology*, 34(1), 101-112.

Cognato, A.I. & Grimaldi, D. (2009) 100 million years of morphological stasis in bark beetles (Coleoptera: Curculionidae: Scolytinae). *Systematic Entomology*, 34, 93–100