

Table 4: Data relating to the presence/absence of faecal pellets beneath tree species used by Koalas in the Bermagui/Murrah Forests. Tree species are grouped in order of the extent to which they satisfied the criteria adopted for analyses purposes.

Tree Species	No.Sites	n	P	Tree Species	No.Sites	n	P
A. Eucalypts				B. Non-eucalypts			
<i>E. agglomerata</i>	30	191	0.073	<i>Acacia cognata</i>	10	22	0.091
<i>E. bosistoana</i>	17	40	0.200	<i>A. falcata</i>	17	37	0.054
<i>E. cypellocarpa</i>	15	41	0.220	<i>A. mearnsii</i>	4	20	0.050
<i>E. globoidea</i>	21	120	0.167	<i>Acmena smithii</i>	6	20	0.100
<i>E. gummifera</i>	24	112	0.071	<i>Allocas. littoralis</i>	52	368	0.087
<i>E. longifolia</i>	62	285	0.214	<i>Ang. floribunda</i>	22	49	0.041
<i>E. muellerana</i>	54	342	0.123	<i>Backhousia myrtifolia</i>	7	71	0.042
<i>E. tricarpa</i>	45	138	0.080	<i>Exo. cupressiformis</i>	7	7	0.143
<i>E. consideniana</i>	6	23	0.304	<i>Acacia implexa</i>	1	1	-
<i>E. maculata</i>	3	41	0.024	<i>A. longifolia</i>	1	1	-
<i>E. seiberi</i>	24	90	0.044	<i>A. irrorata</i>	5	5	-
				<i>Pit. undulatum</i>	1	1	-
<i>E. botryoides</i>	3	5	-	<i>Persoonia linearis</i>	1	1	-
<i>E. elata</i>	1	1	-	<i>Tristaniopsis laurina</i>	1	1	-
<i>E. paniculata</i>	1	2	-				
<i>E. smithii</i>	2	8	-				
Total Trees		1439				604	