<ul> <li>From video, nema= in virtually all parts of earth (soil, salt flats, polar regions, tropics, freshwater, saltwater, hot springs)</li> <li>From video, well over half are:</li> <li>nds in Evolution (Nematoda Characteristics):</li> <li>1. Gastrulation (germ layers):</li> <li>From video, epidermis is called</li> <li>2. Cellular Organization:</li> <li>3. Symmetry:</li> <li>4. Body Cavity: (between mesoderm and endoderm)</li> </ul>	Fast Facts: ●	
From video, nema= in virtually all parts of earth (soil, salt flats, polar regions, tropics, freshwater, saltwater, hot springs)  From video, well over half are:  Inds in Evolution (Nematoda Characteristics):  Gastrulation (gern layers): From video, epidermis is called Cellular Organization: Symmetry: Body Cavity: Getween mesoderm and endoderm)  matoda Body Plan (Essential Roles) Responding to Environment:  In head region (but no "brain") In		
in virtually all parts of earth (soil, salt flats, polar regions, tropics, freshwater, saltwater, hot springs)  From video, well over half are:  Gastrulation (germ layers):  From video, epidermis is called  Cellular Organization:  Symmetry:  Body Cavity:  (between mesoderm and endoderm)  matoda Body Plan (Essential Roles)  Responding to Environment:  Mith  nerve cords  in head region (but no "brain")  to detect chemicals  Movement:  From video,  in body contract and worm  Respiration, Circulation:  From video, nutrients are transported by  Digestion:  1st phyla with  Food enters  Undigested food wits  Excretion:  Anus for undigested food  for metabolic waste  Reproduction:  Most are  (separate male and female)  Males  Parasitic worms have  E.g. Ascaria  Mological Roles of Nematoda  Nematodes excrete the excess nitrogen from their food in plant-available form		
freshwater, saltwater, hot springs)  From video, well over half are:  Inds in Evolution (Nematoda Characteristics):  Cellular Organization:  Symmetry:  Body Cavity:  (between mesoderm and endoderm)  matoda Body Plan (Essential Roles)  Responding to Environment:  in head region (but no "brain")  to detect chemicals  Movement:  From video,  From video,  From video, nutrients are transported by  Digestion:  1st phyla with  Food enters  Undigested food  for metabolic waste  Reproduction:  Most are  (separate male and female)  Fertilization  Males  than females  Parasitic worms have  E.g. Ascaria  Sological Roles of Nematoda  Nematodes excrete the excess nitrogen from their food in plant-available form		
• From video, well over half are:  Inds in Evolution (Nematoda Characteristics):  1. Gastrulation (germ layers):  - From video, epidermis is called  2. Cellular Organization:  3. Symmetry:  4. Body Cavity:  - Body Cavity:  - In head region (but no "brain")  - In hoody contract and worm  - Respiration, Circulation:  - From video, nutrients are transported by  - Digestion:  - 1st phyla with - Food enters - Cours in intestine  - Undigested food exits  - Excretion:  - Anus for undigested food - For metabolic waste  - Reproduction:  - Most are - (separate male and female) - (fertilization - Males - Parasitic worms have - Figerout the process of Nematoda  1. Nematodes excrete the excess nitrogen from their food in plant-available form		pics,
India in Evolution (Nematoda Characteristics):    Gastrulation (germ layers):	· · · · · · · · · · · · · · · · · · ·	
1. Gastrulation (germ layers):  - From video, epidermis is called  2. Cellular Organization:  - Symmetry:  - Body Cavity:  - Movement:  - From video, - in head region (but no "brain") - to detect chemicals  - Movement:  - From video, - in body contract and worm - Respiration, Circulation: - From video, nutrients are transported by - Digestion:  - 1st phyla with - Food enters - Undigested food exits - Excretion: - Anus for undigested food - for metabolic waste  - Reproduction: - Most are - (separate male and female) - Males - Parasitic worms have - Parasitic worms have - Parasitic worms have - E.g. Ascaria    Doligical Roles of Nematoda   Nematodes excrete the excess nitrogen from their food in plant-available form	From video, well over half are:	
1. Gastrulation (germ layers):  - From video, epidermis is called  2. Cellular Organization:  - Symmetry:  - Body Cavity:  - Movement:  - From video, - in head region (but no "brain") - to detect chemicals  - Movement:  - From video, - in body contract and worm - Respiration, Circulation: - From video, nutrients are transported by - Digestion:  - 1st phyla with - Food enters - Undigested food exits - Excretion: - Anus for undigested food - for metabolic waste  - Reproduction: - Most are - (separate male and female) - Males - Parasitic worms have - Parasitic worms have - Parasitic worms have - E.g. Ascaria    Doligical Roles of Nematoda   Nematodes excrete the excess nitrogen from their food in plant-available form		
Symmetry:	·	
2. Cellular Organization: 3. Symmetry: 4. Body Cavity: (between mesoderm and endoderm)  matoda Body Plan (Essential Roles)  • Responding to Environment:		
3. Symmetry:		
### According to Environment:    Responding to Environment:   With	2. Celiulai Organization.	
matoda Body Plan (Essential Roles)  Responding to Environment:    min head region (but no "brain")	Symmetry:      Dedy Cavity:      (between manaderm and endederm)	
Responding to Environment:  with	4. Body Cavity: (between mesodern and endodern)	
with	matoda Body Plan (Essential Roles)	
in head region (but no "brain") to detect chemicals  Movement:	Responding to Environment:	
Movement:  From video, in body contract and worm  Respiration, Circulation:  Through skin, like platyhelminthes  From video, nutrients are transported by  Digestion:  1st phyla with digestive system  Food enters occurs in intestine  Undigested food exits  Excretion:  Anus for undigested food  for metabolic waste  Reproduction:  Most are (separate male and female)  Males than females  Parasitic worms have, like protista and platyhelminthes  E.g. Ascaria  Diogical Roles of Nematoda  Nematodes excrete the excess nitrogen from their food in plant-available form	o with ner	rve cords
Movement:		
<ul> <li>From video, in body contract and worm</li></ul>	o to detect chemicals	
Respiration, Circulation:  through skin, like platyhelminthes  From video, nutrients are transported by  Digestion:  1st phyla with digestive system Food enters occurs in intestine  Undigested food exits  Excretion:  Anus for undigested food for metabolic waste  Reproduction:  Most are device ring excretory pore excretory	Movement:	
through skin, like platyhelminthes  From video, nutrients are transported by  Digestion:  1st phyla with digestive system  Food enters occurs in intestine  Undigested food exits  Excretion:  Anus for undigested food  for metabolic waste  Reproduction:  Most are (separate male and female)  Males than females  Parasitic worms have , like protista and platyhelminthes  E.g. Ascaria  Diogical Roles of Nematoda  Nematodes excrete the excess nitrogen from their food in plant-available form  ()	<ul> <li>From video, in body contract and worm</li> </ul>	
Digestion:   Other properties are transported by	·	
Digestion:  1st phyla with	<ul> <li>through skin, like platyhelminthes</li> </ul>	
o 1st phyla with		
Food enters occurs in intestine      Undigested food exits      Excretion:      Anus for undigested food mouth intestine psuedocoleom ocuticle for metabolic waste      Reproduction:      Most are device ring excretory pore ovary reproductive pore ovary reproductive pore in the pharynx ovary reproductive pore ovary ovary reproductive pore ovary pore ova	•	
occurs in intestine Undigested food exits  Anus for undigested food for metabolic waste  Reproduction:  Most are		
<ul> <li>Undigested food exits</li></ul>		
Excretion:		
Anus for undigested food  for metabolic waste  Reproduction:  Most are	<u> </u>	
Anus for undigested food  for metabolic waste  Reproduction:  Most are fertilization  Males than females  Parasitic worms have, like protista and platyhelminthes  E.g. Ascaria  Diogical Roles of Nematoda  Nematodes excrete the excess nitrogen from their food in plant-available form  ()	mouth	
for metabolic waste  Reproduction:  Most are	Anus for undigested food	
Reproduction:  Most are		
Most are device ring excretory pore ovary reproductive pore (separate male and female)  Males than females Parasitic worms have, like protista and platyhelminthes E.g. Ascaria  Nematodes excrete the excess nitrogen from their food in plant-available form  ()		
<ul> <li>Most are piercing device ring</li> <li>Most are piercing device ring</li> <li>Most are piercing nerve reproductive pore</li> <li>(separate male and female)</li> <li>Males than females</li> <li>Parasitic worms have, like protista and platyhelminthes</li> <li>E.g. Ascaria</li> <li>Nematodes of Nematoda</li> <li>Nematodes excrete the excess nitrogen from their food in plant-available form ()</li> </ul>	retracted	
<ul> <li>Most are tevice ring</li></ul>	piercing page	reproductiv
<ul> <li> fertilization</li> <li>Males than females</li> <li>Parasitic worms have, like protista and platyhelminthes</li> <li>E.g. Ascaria</li> </ul> Diogical Roles of Nematoda <ol> <li>Nematodes excrete the excess nitrogen from their food in plant-available form</li> <li>)</li> </ol>	Most are ring	•
<ul> <li>Males than females</li> <li>Parasitic worms have, like protista and platyhelminthes</li> <li>E.g. Ascaria</li> </ul> <b>blogical Roles of Nematoda</b> <ol> <li>Nematodes excrete the excess nitrogen from their food in plant-available form</li> <li>)</li> </ol>		
<ul> <li>Parasitic worms have, like protista and platyhelminthes</li> <li>E.g. Ascaria</li> <li>Plogical Roles of Nematoda</li> <li>Nematodes excrete the excess nitrogen from their food in plant-available form ()</li> </ul>		
<ul> <li>E.g. Ascaria</li> <li>blogical Roles of Nematoda</li> <li>Nematodes excrete the excess nitrogen from their food in plant-available form         <ul> <li>()</li> </ul> </li> </ul>		41
Diogical Roles of Nematoda  1. Nematodes excrete the excess nitrogen from their food in plant-available form  ()		ines
Nematodes excrete the excess nitrogen from their food in plant-available form  ()	∪ E.g. Ascana	
Nematodes excrete the excess nitrogen from their food in plant-available form  ()	ological Polog of Namatoda	
()		
2 Recause of their abundance, diversity and effects on soil process indices of nematode assemblages	veae. exe.e .ue ex. essuu.uueu uuuu uueu uuuu 1177711-3773113019 101111	
Z. Because of mell andinance, inversity and energy of coil biorded indicae of namatoda decombidade.	1. We materials a character the choose mitrogen from their rood in plant available form	