

## References

Rassu, F.S., McFadden, M., Aaron, R.V., Wegener, S.T., Ephraim, P.L, Lane, E., Brennan, G., Minick, K.I., Fritz, J.M., Skolasky, R.L. (2021, June) The Impact Of The COVID-19 Pandemic on The Relationship Between Neighborhood Deprivation And Pain-Related Experiences Among U.S. Patients With Chronic Low Back Pain. Poster presented at the International Association for the Study of Pain (IASP) World Congress of Pain (Online Conference)

1. Skolasky, R. L. et al. The OPTIMIZE study: protocol of a pragmatic sequential multiple assessment randomized trial of nonpharmacologic treatment for chronic, nonspecific low back pain. *BMC Musculoskelet. Disord.* 1–14 (2020).
2. Fischer, D. et al. Capturing the patient's view of change as a clinical outcome measure. *J. Am. Med. Assoc.* 282, 1157–1162 (1999).
3. Kind, A. J. H. & Buckingham, W. R. Making Neighborhood-Disadvantage Metrics Accessible — The Neighborhood Atlas. *N. Engl. J. Med.* 378, 2456–2458 (2018).
4. University of Wisconsin School of Medicine and Public Health. Area Deprivation Index v3.0. <https://www.neighborhoodatlas.medicine.wisc.edu/> (2018).
5. Cella, D. et al. The patient-reported outcomes measurement information system (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005-2008. *J. Clin. Epidemiol.* 63, 1179–1194 (2010).