Feeding Intervention
 5 point wipe (wipe out to in to eye, mouth, chin firm pressure to desensitize) Vibrating toothbrush on oral cavity. (move it around gums and mouth for 5 minutes to warm up mouth) Z vibe on cheeks, roof of mouth, tongue, gums (rubber vibrate to desensitize, 5 min) Firm pressure inside of mouth (child directed) [pushing up on hard palate, desensitize) Nuk brush on outer teeth and gums (child directed)
 5 point wipe with vibration Proprioceptive input in mouth (chewing gum, sucking/blowing activities with whistle toys, blow bubbles, blow in straw to make bubbles, drink from small straw to get tight lips, blow cotton ball with straw) [to use muscles of mouth and purse their lips over something] Bandana scarf to protect clothes and skin (rashy)
 Long shaped teethers (teether in cheek, the tongue automatically moves toward it) [important for moving food around mouth] Nuk with peanut butter on side of mouth [they will need to lick peanut butter off with tonge] Use different textured foods. [move the food around in the mouth] Cheese cloth or chew bags on side of

	5. When beginning chewing-use easily dissolvable food {texture they can easily chew and tolerate}
Abnormal Bite/tonic bit	 Maroon Spoon (avoid plastic spoons) {don't want to break plastic spoon off and break teeth with metal spoon, they need to do chin tuck, jaw stimulation to try to loosen their jaw hold on spoon} Present food below eye level to encourage chin tuck. Avoid scraping food on teeth. {at eye level or a little below, don't scrape their teeth and lips on food bc it can elicit tonic bite (narrow bowl on spoon)} If bit elicited-wait for mouth to open.
Abnormal Gag Reflex	 Relaxed, unhurried atmosphere Tip head forward if gag initiated Downward pressure on tongue with spoon.
Drinking	Honeybear for straw use- squeeze honey bear (to teach them to start using straw)

honeybear for tongue thrust