

AMUKURA PARISH JOINT EXAM 2021

CHEM PP2 MARKING SCHEME

- 1.(i) Noble gases $\sqrt{1}$
- (ii) D_2SO_4 $\sqrt{1}$
- (iii) (a) Y $\sqrt{1}$
(b) E $\sqrt{1}$
- (iv) Ionic bond $\sqrt{1}$ – Because B reacts by losing an electron (s) which are gained by H.
 $\sqrt{1}$ accept transfer of electrons from a metal to non metal
- (v) D//M $\sqrt{1}$ Any $\frac{1}{2}$ mark each
- (vi) Because E reacts by gaining an extra electron which reduces $\sqrt{1}$ the electrostatic pull by the positive nucleus making the ionic radius increase. Or incoming electron causes increased repulsion wtte
- (vii) At Period III Group IV
- (viii) Because of the increase in the strength of the molecular bonds in the oxide of L as compared to that of G. $\sqrt{1}$ w.t.t.e
- (ix) C has a smaller atomic $\sqrt{1}$ radius than I because of the increase in the strength of the
Nuclear force of attraction in C as the number of protons increase $\sqrt{1}$ w.t.t.e
- (x) 1st ionization energies increases from J – L across the period due to addition of an extra proton in the nucleus increasing the attraction of the valency electrons $\sqrt{1}$
- 2 a) i) A solution that cannot dissolve any more of the solute at that particular temperature. ✓
1mk
- ii) Scientific technique used to separate substances due to their differences in their crystallization temperature. ✓ 1mk or w.t.t.e
- b) i) on the scanned graph
- ii) $x=100g/100ml$, $y=40g/100ml$

iii) 5°C

iv) type of hardness that cannot be removed by boiling

3. a) (i) Fractional distillation ✓ 1mk

(ii) Argon//neon/xenon//krypton ✓ 1mk

- b)
- | | | |
|---|--------------------|-------|
| A | Sulphur | ✓ 1mk |
| B | Ammonia gas | ✓ 1mk |
| C | sulphur (vi) oxide | ✓ 1mk |
| D | Ammonium sulphate | ✓ 1mk |

c) (i) Finely divided iron ✓ 1mk

(ii) Vanadium (v) oxide ✓ 1mk

(iii) The catalysts fasten ✓ 1mk the Haber & contact processes by lowering the activation energy ✓ 1mk of the reactions//the rate of production is increased.

d) (i) $\text{H}_2\text{SO}_{4(\text{aq})} + 2\text{NH}_{3(\text{g})} \rightarrow (\text{NH}_4)_2\text{SO}_{4(\text{aq})}$ ✓ 1mk

(ii) Formula mass of $(\text{NH}_4)_2\text{SO}_4 = 2(14+4) + 32 + 4(16)$

$= 132 \text{ grams}$ ✓ ½ mk


$$\% \text{ of N} = \frac{28}{132} \times 100 \quad \checkmark 1\text{mk}$$
$$= 21.212\% \quad \checkmark \frac{1}{2} \text{mk}$$

(iii) Use as a fertilizer ✓ 1mk

4. a) I: The outlet delivery tube should not dip into the Zinc/dilute Sulphuric acid mixture in the round bottomed flask. ✓ 1mk

II: The use of heat is not required ✓ 1mk

b)



Conc H_2SO_4

1mk

c) i) It is denser than air ✓ 1 mk

ii) $\text{H}_{2(\text{g})} + \frac{1}{2} \text{O}_{2(\text{g})} \rightarrow \text{H}_2\text{O}_{(\text{g})}$ ✓ balancing $\frac{1}{2}$ mark
states $\frac{1}{2}$ mark

d) $\text{Zn}_{2(\text{s})} + \text{H}_2\text{SO}_{4(\text{aq})} \rightarrow \text{ZnSO}_{4(\text{aq})} + \text{H}_{2(\text{g})}$ balancing $\frac{1}{2}$ mk
states $\frac{1}{2}$ mk

$$\begin{array}{ccc} 1 \text{ vol} & 1 \text{ vol} & 1 \text{ vol} \\ \left[\frac{6.54}{R} \right] & & \left[\frac{2.4}{24} \right] \end{array}$$

Therefore, $\left[\frac{6.54}{R} \right] = \frac{2.4}{24}$, ✓ 1mk where $R = \text{R.A.M of Zinc}$

$$R = \frac{24 \times 6.54}{2.4}$$

Or $R = 65.4$ ✓ 1mk

e) - $\text{H}_{2(\text{g})}$ is used in balloons by meteorologists ✓ 1mk

- It is used as rocket fuel ✓ 1mk

5.

(a) Heating copper (ii) oxide ✓ 1mk

(b) Black solid would turn brown ✓ 1mk

(c) $\text{CuO}_{(\text{s})} + \text{CO}_{(\text{g})} \rightarrow \text{Cu}_{(\text{s})} + \text{CO}_{2(\text{g})}$ ✓ 1 $\frac{1}{2}$ mk

(d) $2\text{CO}_{(\text{g})} + \text{O}_{2(\text{g})} \rightarrow 2\text{CO}_{2(\text{g})}$ ✓ 1 $\frac{1}{2}$ mk

(e) It is poisonous ✓ 1mk

(f) (i) Reducing agent - Carbon(ii) oxide ✓ 1mk

(ii) Oxidising agent - Copper (ii) oxide ✓ 1mk

(g) Hydrogen / ammonia gas (Any one) ✓ 1mk

(h) There would be no observable change ✓ 1mk. This is because sodium is higher than carbon in the reactivity series and therefore has higher affinity of oxygen ✓ 1mk

6.

a) Concentrated sulphuric (vi) acid ✓ 1mk

b) It is denser than air ✓ 1mk

c) It turns red then white. ✓ 1mk

It turns white / it gets bleached ✓ 1mk

d) $\text{Cl}_{2(\text{g})} + \text{H}_2\text{O}_{(\text{l})} \rightarrow \text{HOCl}_{(\text{aq})} + \text{HCl}_{(\text{aq})}$ ✓ 1mk

e) PCl_3 ✓ 1mk

PCl_5 $\sqrt{1mk}$

- f) A yellow deposit of sulphur is formed / seen $\sqrt{1mk}$
Chlorine oxidizes sulphide ions to solid sulphur $\sqrt{1mk}$

g)

- Manufacture of hydrochloric acid $\sqrt{1mk}$
- Manufacture of bleaching agents such as chlorate used in the cotton and paper industries
- Chlorine is used in the treatment of water and sewage plants
- Manufacture of chloroform as an anaesthetic
- Manufacture of solvents such as trichloroethane

Any one

7.

- a) A - Filtration $\sqrt{1 \frac{1}{2} mk}$
B - Absorption $\sqrt{1 \frac{1}{2} mk}$
M - Isolation of water $\sqrt{1 \frac{1}{2} mk}$
D - Cooling $\sqrt{1 \frac{1}{2} mk}$
- b) Liquids – NaOH (aq) / KOH (aq) $\sqrt{1mk}$
Substance T – Ice / water $\sqrt{1mk}$
- c) To increase surface area for cooling $\sqrt{1 mk}$
- d) (i) Oxygen is used to remove impurities during steel making $\sqrt{1 mk}$
(ii) Is used in cutting and welding of metals $\sqrt{1 mk}$
- e) $2\text{H}_2\text{O}_{2(l)} \xrightarrow{\text{MnO}_{2(s)}} 2\text{H}_2\text{O}_{(l)} + \text{O}_{2(g)}$ $\sqrt{1 mk}$
- f) (i) R - Rusting occurred $\sqrt{1 \frac{1}{2} mk}$ because of air and water being present $\sqrt{\frac{1}{2} mk}$
S - No rusting $\sqrt{\frac{1}{2} mk}$ Water is absent $\sqrt{\frac{1}{2} mk}$
T - No rusting $\sqrt{\frac{1}{2} mk}$ Air is absent $\sqrt{\frac{1}{2} mk}$
(ii) To prevent rusting $\sqrt{1mk}$
To increase aesthetic value of the metal

