

References

- Bartsch, T., Döhring, J., Reuter, S., Finke, C., Rohr, A., Brauer, H., Deuschl, G., & Jansen, O. (2015). Selective neuronal vulnerability of human hippocampal CA1 neurons: lesion evolution, temporal course, and pattern of hippocampal damage in diffusion-weighted MR imaging. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*, 35(11), 1836–1845. <https://doi.org/10.1038/jcbfm.2015.137>
- Bartsch, T., Döhring, J., Rohr, A., Jansen, O., & Deuschl, G. (2011). CA1 neurons in the human hippocampus are critical for autobiographical memory, mental time travel, and autonoetic consciousness. *Proceedings of the National Academy of Sciences of the United States of America*, 108(42), 17562–17567. <https://doi.org/10.1073/pnas.1110266108>
- Dimitrova, L. I., Dean, S. L., Schlumpf, Y. R., Vissia, E. M., Nijenhuis, E. R. S., Chatzi, V., Jäncke, L., Veltman, D. J., Chalavi, S., & Reinders, A. A. T. S. (2023). A neurostructural biomarker of dissociative amnesia: a hippocampal study in dissociative identity disorder. *Psychological medicine*, 53(3), 1–9. Advance online publication. <https://doi.org/10.1017/S0033291721002154>
- Ehling, T., Nijenhuis, E. R., & Krikke, A. P. (2008). Volume of discrete brain structures in complex dissociative disorders: preliminary findings. *Progress in brain research*, 167, 307–310. [https://doi.org/10.1016/S0079-6123\(07\)67029-0](https://doi.org/10.1016/S0079-6123(07)67029-0)
- Forrest K. A. (2001). Toward an etiology of dissociative identity disorder: a neurodevelopmental approach. *Consciousness and cognition*, 10(3), 259–293. <https://doi.org/10.1006/ccog.2001.0493>
- Semple, W. E., Goyer, P. F., McCormick, R., Donovan, B., Muzic, R. F., Jr, Rugle, L., McCutcheon, K., Lewis, C., Liebling, D., Kowaliw, S., Vapenik, K., Semple, M. A., Flener, C. R., & Schulz, S. C. (2000). Higher brain blood flow at amygdala and lower frontal cortex blood flow in PTSD patients with comorbid cocaine and alcohol abuse compared with normals. *Psychiatry*, 63(1), 65–74. <https://doi.org/10.1080/00332747.2000.11024895>
- Shin, L. M., Whalen, P. J., Pitman, R. K., Bush, G., Macklin, M. L., Lasko, N. B., Orr, S. P., McInerney, S. C., & Rauch, S. L. (2001). An fMRI study of anterior cingulate function in posttraumatic stress disorder. *Biological psychiatry*, 50(12), 932–942. [https://doi.org/10.1016/s0006-3223\(01\)01215-x](https://doi.org/10.1016/s0006-3223(01)01215-x)
- Shin, L. M., Rauch, S. L., & Pitman, R. K. (2006). Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Annals of the New York Academy of Sciences*, 1071(1), 67-79. <https://doi.org/10.1196/annals.1364.007>
- Vermetten, E., Schmahl, C., Lindner, S., Loewenstein, R. J., & Bremner, J. D. (2006). Hippocampal and amygdalar volumes in dissociative identity disorder. *The American journal of psychiatry*, 163(4), 630–636. <https://doi.org/10.1176/ajp.2006.163.4.630>