

[See this page in the course material.](#)

Learning Outcomes

- Simplify n th roots.
- Write radicals as rational exponents.

Launch Desmos Calculator

Using Rational Roots

Although square roots are the most common rational roots, we can also find cube roots, 4th roots, 5th roots, and more. Just as the square root function is the inverse of the squaring function, these roots are the inverse of their respective power functions. These functions can be useful when we need to determine the number that, when raised to a certain power, gives a certain number.

Suppose we know that $a^3 = 8$. We want to find what number raised to the 3rd power is equal to 8. Since $2^3 = 8$, we say that 2 is the cube root of 8.

The n th root of a is a number that, when raised to the n th power, gives a . For example, -3 is the 5th root of -243 because $(-3)^5 = -243$. If a is a real number with at least one n th root, then the **principal n th root** of a is the number with the same sign as a that, when raised to the n th power, equals a .

The principal n th root of a is written as $\sqrt[n]{a}$, where n is a positive integer greater than or equal to 2. In the radical expression, n is called the **index** of the radical.

A General Note: Principal n th Root

If a is a real number with at least one n th root, then the **principal n th root** of a , written as $\sqrt[n]{a}$, is the number with the same sign as a that, when raised to the n th power, equals a . The **index** of the radical is n .

Simplify each of the following:

Simplify each of the following:

- Show Solution

- ## Try It

Simplify.

- Show Solution

1. -6
2. 6
3. $88\sqrt[3]{9}$



[See this interactive in the course material.](#)



[See this interactive in the course material.](#)



[See this interactive in the course material.](#)



[See this interactive in the course material.](#)

Using Rational Exponents

Radical expressions can also be written without using the radical symbol. We can use rational (fractional) exponents. The index must be a positive integer. If the index n is even, then a cannot be negative.

$$[latex]{a}^{\frac{1}{n}}=\sqrt[n]{a}[/latex]$$

We can also have rational exponents with numerators other than 1. In these cases, the exponent must be a fraction in lowest terms. We raise the base to a power and take an n th root. The numerator tells us the power and the denominator tells us the root.

$$[latex]{a}^{\frac{m}{n}}=\left(\sqrt[n]{a}\right)^m=\sqrt[n]{{a}^m}[/latex]$$

All of the properties of exponents that we learned for integer exponents also hold for rational exponents.

Rational Exponents

Rational exponents are another way to express principal n th roots. The general form for converting between a radical expression with a radical symbol and one with a rational exponent is

$$[latex]\begin{align}{a}^{\frac{m}{n}}&=\left(\sqrt[n]{a}\right)^m=\sqrt[n]{{a}^m}\end{align}[/latex]$$

How To: Given an expression with a rational exponent, write the expression as a radical.

1. Determine the power by looking at the numerator of the exponent.
2. Determine the root by looking at the denominator of the exponent.
3. Using the base as the radicand, raise the radicand to the power and use the root as the index.

Example: Writing Rational Exponents as Radicals

Write $[latex]{343}^{\frac{2}{3}}[/latex]$ as a radical. Simplify.

Show Solution

The 2 tells us the power and the 3 tells us the root.

$$[latex]{343}^{\frac{2}{3}}=\left(\sqrt[3]{343}\right)^2=\sqrt[3]{{343}^2}[/latex]$$

We know that $\sqrt[3]{343}=7$ because $7^3=343$. Because the cube root is easy to find, it is easiest to find the cube root before squaring for this problem. In general, it is easier to find the root first and then raise it to a power.

$$343^{\frac{2}{3}} = (\sqrt[3]{343})^2 = 7^2 = 49$$

Try It

Write $9^{\frac{5}{2}}$ as a radical. Simplify.

Show Solution

$$(\sqrt{9})^5 = 3^5 = 243$$



[See this interactive in the course material.](#)

Example: Writing Radicals as Rational Exponents

Write $\frac{4}{\sqrt[7]{a^2}}$ using a rational exponent.

Show Solution

The power is 2 and the root is 7, so the rational exponent will be $\frac{2}{7}$. We get $a^{\frac{4}{\sqrt{7}}}$. Using properties of exponents, we get $a^{\frac{4}{\sqrt{7}} \cdot \frac{1}{\sqrt{7}}} = a^{\frac{4}{7}}$.

Try It

Write $x\sqrt[9]{5y}$ using a rational exponent.

Show Solution

$$x(5y)^{\frac{1}{9}}$$

Watch this video to see more examples of how to write a radical with a fractional exponent.



[Video Link](#)

Example: Simplifying Rational Exponents

Simplify:

- $5\left(2x^{\frac{3}{4}}\right)\left(3x^{\frac{1}{5}}\right)$
- $\left(\frac{16}{9}\right)^{-\frac{1}{2}}$

Show Solution

1.

```
[latex]\begin{align}&30{x}^{\frac{3}{4}}{x}^{\frac{1}{5}}&& \text{Multiply the coefficients}. \\&30{x}^{\frac{3}{4}+\frac{1}{5}}&& \text{Use properties of exponents}. \\&30{x}^{\frac{19}{20}}&& \text{Simplify}. \end{align}
```

2.

```
[latex]\begin{align}&\left(\frac{9}{16}\right)^{\frac{1}{2}}&& \text{Use definition of negative exponents}. \\&\sqrt{\frac{9}{16}}&& \text{Rewrite as a radical}. \\&\frac{\sqrt{9}}{\sqrt{16}}&& \text{Use the quotient rule}. \\&\frac{3}{4}&& \text{Simplify}. \end{align}
```

Try It

Simplify
$$\left(8x\right)^{\frac{1}{3}}\left(14x^{\frac{6}{5}}\right)^{\frac{1}{2}}$$

Show Solution

```
[latex]28{x}^{\frac{23}{15}}
```



[See this interactive in the course material.](#)

Licenses and Attributions

CC licensed content, Original

- Revision and Adaptation. **Provided by:** Lumen Learning. **License:** [CC BY: Attribution](#)

CC licensed content, Shared previously

- College Algebra. **Authored by:** Abramson, Jay et al.. **Provided by:** OpenStax. **Located at:** <http://cnx.org/contents/9b08c294-057f-4201-9f48-5d6ad992740d@5.2>. **License:** [CC BY: Attribution](#). **License Terms:** Download for free at <http://cnx.org/contents/9b08c294-057f-4201-9f48-5d6ad992740d@5.2>
- Ex: Write a Radical in Rational Exponent Form. **Authored by:** James Sousa (Mathispower4u.com). **Located at:** https://youtu.be/L5Z_3RrVjA. **License:** [CC BY: Attribution](#)
- Question ID 2564, 2565, 2567, 2592. **Authored by:** Greg Langkamp. **License:** [CC BY: Attribution](#). **License Terms:** IMathAS Community License, CC-BY + GPL
- Question ID 3415. **Authored by:** Jessica Reidel. **License:** [CC BY: Attribution](#). **License Terms:** IMathAS Community License, CC-BY + GPL
- Question ID 59783. **Authored by:** Gary Parker. **License:** [CC BY: Attribution](#). **License Terms:** IMathAS Community License, CC-BY + GPL

CC licensed content, Specific attribution

- College Algebra. **Authored by:** OpenStax College Algebra. **Provided by:** OpenStax. **Located at:** <http://cnx.org/contents/9b08c294-057f-4201-9f48-5d6ad992740d@3.278:1/Preface>. **License:** [CC BY: Attribution](#)

</div